Recombinant human CRLF2 protein

Catalog Number: ATGP3478

PRODUCT INFORMATION

Expression system
Baculovirus

Domain
23-231aa

UniProt No.
Q9HC73

NCBI Accession No.
NP_071431.2

Alternative Names
Cytokine receptor-like factor 2 isoform 1 precursor, CRL2, CRLF2Y, TSLPR, Cytokine receptor-like factor 2 isoform 1, CRLF2, Cytokine receptor-like 2, ILXR, Thymic stromal lymphopoietin protein receptor, TSLP receptor

PRODUCT SPECIFICATION

Molecular Weight
25.2 kDa (218aa)

Concentration
1mg/ml (determined by absorbance at 280nm)

Formulation
Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

Purity
> 95% by SDS-PAGE

Endotoxin level
< 1 EU per 1ug of protein (determined by LAL method)

Tag
His-Tag

Application
SDS-PAGE

Storage Condition
Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description
CRLF2, also known as cytokine receptor-like factor 2 isoform 1, is a receptor for thymic stromal lymphopoietin. It is proposed to signal through a heterodimeric receptor complex that consists of a new member of the hemopoietin family termed CRLF2 and the IL-7R alpha-chain. The protein induces phosphorylation of Stat3 and

Recombinant human CRLF2 protein

Catalog Number: ATGP3478

Stat5 also activate JAK2 pathways. CRLF2 and IL-7R alpha are principally coexpressed on monocytes and dendritic cell populations and to a much lesser extent of various lymphoid cells. Recombinant human CRLF2, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

Amino acid Sequence

<ADP>QGGAAEG VQIQIIYFNL ETVQVTWNAS KYSRTNLTFH YRFNGDEAYD QCTNYLLQEG HTSGCLLDAE QRDDILYFSI RNGTHPVFTA SRWMVYYLKP SSPKHVRFSW HQDAVTVTCS DLSYGDLLYE VQYRSPFDTE WQSKQENTCN VTIEGLDAEK CYSFWVRVKA MEDVYGPDTY PSDWSEVTCW QRGEIRDACA ETPTPPKPKL SK<HHHHHH>

General References

Reche PA., et al. (2001) J Immunol. 167:336-343.
Chen IM., et al. (2012) Blood. 119:3512-3522.

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain