# **PRODUCT INFORMATION**

**Expression system** Baculovirus

**Domain** 21-157aa

**UniProt No.** Q99969

NCBI Accession No. NP\_002880

Alternative Names Retinoic acid receptor responder protein 2, RARRES2, HP10433, TIG2, Chemerin

# **PRODUCT SPECIFICATION**

Molecular Weight 16.9 kDa (146aa)

**Concentration** 0.5mg/ml (determined by absorbance at 280nm)

### Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

### Purity

> 90% by SDS-PAGE

**Endotoxin level** < 1 EU per 1ug of protein (determined by LAL method)

**Tag** His-Tag

Application SDS-PAGE

### **Storage Condition**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

### BACKGROUND

### Description

RARRES2, as known as retinoic acid receptor responder protein 2, is a distant member of the Cystatin superfamily. Members of this superfamily contain at least two intra-chain disulfide bonds and an alpha-helical structure over a distance of about 100 amino acids. This protein was found to stimulate chemotaxis of dendritic cells and macrophages to the site of inflammation. Also, it has been implicated in autocrine/paracrine signaling



for adipocyte differentiation and also stimulation of lipolysis. Recombinant human RARRES2, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

#### **Amino acid Sequence**

ADPELTEAQR RGLQVALEEF HKHPPVQWAF QETSVESAVD TPFPAGIFVR LEFKLQQTSC RKRDWKKPEC KVRPNGRKRK CLACIKLGSE DKVLGRLVHC PIETQVLREA EEHQETQCLR VQRAGEDPHS FYFPGQFAFS HHHHHH

### **General References**

Hatziagelaki E., et al, (2015) PLoS ONE 10:E0124935. Zabel BA., et al, (2014) Am J Clin Exp Immunol. 3:1-19.

## DATA



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

15% SDS-PAGE (3ug)