PRODUCT INFORMATION

Expression system Baculovirus

Domain 19-328aa

UniProt No. Q01638

NCBI Accession No. NP_057316.3

Alternative Names

Interleukin-1 receptor-like 1, Protein ST2, homolog of mouse growth stimulation-expressed, DER4, T1, FIT-1, ST2L, ST2V, IL33R

PRODUCT SPECIFICATION

Molecular Weight

36 kDa (318aa)

Concentration

0.5mg/ml (determined by absorbance at 280nm)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

Purity > 95% by SDS-PAGE

Endotoxin level

< 1 EU per 1ug of protein (determined by LAL method)

Biological Activity

Measured by its ability to inhibit proliferation using D10.G4.1 mouse helper T cells. The ED50 range \leq 10ng/ml with Human IL-33.

Tag His-Ta

His-Tag

Application SDS-PAGE, Bioactivity

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.



BACKGROUND

Description

IL1RL1, as known as interleukin-1 receptor-like 1 isoform 1, is and interleukin-1 receptor family glycoprotein that contributes to immune responses. As an important mediator involved in many immune and inflammatory responses, this cytokine has been implicated as a regulator of both the development and effector phases of type 2 helper T cell responses, and as a negative feedback modulator of macrophage pro-inflammatory function. Recombinant human IL1RL1, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

Amino acid Sequence

KFSKQSWGLE NEALIVRCPR QGKPSYTVDW YYSQTNKSIP TQERNRVFAS GQLLKFLPAA VADSGIYTCI VRSPTFNRTG YANVTIYKKQ SDCNVPDYLM YSTVSGSEKN SKIYCPTIDL YNWTAPLEWF KNCQALQGSR YRAHKSFLVI DNVMTEDAGD YTCKFIHNEN GANYSVTATR SFTVKDEQGF SLFPVIGAPA QNEIKEVEIG KNANLTCSAC FGKGTQFLAA VLWQLNGTKI TDFGEPRIQQ EEGQNQSFSN GLACLDMVLR IADVKEEDLL LQYDCLALNL HGLRRHTVRL SRKNPIDHHS <LEHHHHHH>

General References

Yu XX., et al. (2015) Dig. Dis. Sci. 60:1265-1272. Rolland T., et al. (2014) Int. J. Mol. Sci. 15:23227-23239.

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.





Measured by its ability to inhibit proliferation using D10.G4.1 mouse helper T cells. The ED50 range \leq 10 ng/ml with Human IL-33.