NKMAXBio We support you, we believe in your research

Recombinant canine IL-1 alpha/IL1A protein

Catalog Number: ATGP3976

PRODUCT INFORMATION

Expression system

Baculovirus

Domain

109-265aa

UniProt No.

046612

NCBI Accession No.

NP 001003157

Alternative Names

IL-1 alpha, Il-1a, Interleukin 1 alpha, Interleukin-1 alpha precursor

PRODUCT SPECIFICATION

Molecular Weight

19.3kDa (166aa)

Concentration

0.25mg/ml (determined by Absorbance at 280nm)

Formulation

Liquid. In Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

Purity

> 90% by SDS - PAGE

Endotoxin level

< 1 EU per 1ug of protein (determined by LAL method)

Biological Activity

Measured in a cell proliferation assay using D10.G4.1 mouse helper T cell. The ED50 range ≤ 2 ng/ml.

Tag

His-Tag

Application

SDS-PAGE, Bioactivity

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

IL-1 alpha /IL-1F1, also known as hematopoietin 1 is a cytokine of the interleukin 1 family. IL-1 α is a unique



NKMAXBIO We support you, we believe in your research

Recombinant canine IL-1 alpha/IL1A protein

Catalog Number: ATGP3976

member in the cytokine family in the sense that the structure of its initially synthesized precursor does not contain a signal peptide fragment. In general, Interleukin 1 is responsible for the production of inflammation, as well as the promotion of fever and sepsis. IL-1 alpha inhibitors are being developed to interrupt those processes and treat diseases. IL-1 alpha is produced mainly by activated macrophages, as well as neutrophils, epithelial cells, and endothelial cells. It possesses metabolic, physiological, haematopoietic activities, and plays one of the central roles in the regulation of the immune responses. Although IL-1 production is generally considered to be a consequence of inflammation, recent evidence suggests that IL-1 is also temporarily upregulated during bone formation and the menstrual cycle and can be induced in response to nervous system stimulation. Recombinant canine IL-1 alpha /IL-1F1, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

Amino acid Sequence


<ADP>SVAYNFH NNEKYNYIRI IKSQFILNDN LNQSIVRQTG GNYLMTAALQ NLDDAVKFDM GAYTSEDSKL PVTLRISKTR LFVSAQNEDE PVLLKEMPET PKTIRDETNL LFFWERHGSK HYFKSVAQPK LFIATQERKL VHMARGQPSI TDFRLLETQP < HHHHHHH>

General References

Watanabe N., et al, (1994) Cytokine. 6:597-601. Nicklin MJ., et al, (1994) Genomics. 19:382-384. March CJ, Mosley B., et al, (1985). Nature. 315: 641-647.

DATA

SDS-PAGE




3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain

Biological Activity



Recombinant canine IL-1 alpha/IL1A protein

Catalog Number: ATGP3976



Canine IL-1 alpha/IL-1F1 in a cell proliferation assay using D10.G4.1 mouse helper T cell. The ED50 range \leq 2 ng/ml.

