

Recombinant human FAP protein

Catalog Number: ATGP3988

PRODUCT INFORMATION

Expression system

Baculovirus

Domain

26-760aa

UniProt No.

Q12884

NCBI Accession No.

NP_004451.2

Alternative Names

Fibroblast activation protein, DPPIV Protein, DPPIV, FAPA, Fapalpha Protein, SIMP Protein, Prolyl endopeptidase FAP, 170 kDa melanoma membrane-bound gelatinase, Dipeptidyl peptidase FAP, Fibroblast activation protein alpha, FAPalpha, Gelatine degradation protease FAP, Integral membrane serine protease, Post-proline cleaving enzyme, Serine integral membrane protease, SIMP, Surface-expressed protease, Seprase

PRODUCT SPECIFICATION

Molecular Weight

86.1 kDa (744aa)

Concentration

0.25mg/ml (determined by absorbance at 280nm)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 20% glycerol

Purity

> 90% by SDS-PAGE

Endotoxin level

< 1 EU per 1ug of protein (determined by LAL method)

Biological Activity

Specific activity is > 5,000 pmol/min/ug, and is defined as the amount of enzyme that hydrolysis 1.0 pmole of Z-GP-AMC per minute at pH 7.5, at 37C.

Tag

His-Tag

Application

SDS-PAGE, Enzyme Activity

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

Recombinant human FAP protein

Catalog Number: ATGP3988

BACKGROUND

Description

FAP, also known as Seprase, is a homodimeric integral membrane gelatinase belonging to the serine protease family. FAP includes dipeptidyl peptidase IV (DPPIV / CD26) and related type II transmembrane prolyl serine peptidases, which exert their mechanisms of action on the cell surface. And its enzymatic activity is dependent on FAP association with DPPIV on the cell surface. The active site of FAP is localized in the extracellular part of the protein and contains a catalytic triad composed of Ser624 Asp702 His734 in humans and mice. It is catalytically active as a 170kD homodimer and has a dipeptidase and an endopeptidase activity. FAP expression is high in reactive stromal fibroblasts of epithelial cancers, granulation tissue of healing wounds, and malignant cells of bone and soft tissue sarcomas. FAP is thought to be involved in the control of fibroblast growth or epithelial-mesenchymal interactions during development, tissue repair, and epithelial carcinogenesis. recombinant human FAP, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

Amino acid Sequence


<ADP>LRPSRVH NSEENTMRAL TLKDILNGTF SYKTFFPNWI SGQEYLHQSA DNNIVLYNIE TGQSYTILSN RTMKSVNASN YGLSPDRQFV YLESDYSKLW RYSYTATYYI YDLSNGEFVR GNELPRPIQY LCWSPVGSKL AYVYQNNIYL KQRPDPPFQ ITFNGRENKI FNGIPDWVYE EEMLATKYAL WWSPNGKFLA YAEFNDTDIP VIAYSYYGDE QYPTINIPY PKAGAKNPVV RIFIIDTTYP AYVGPQEVVP PAMIASSDYY FSWLTWVTDE RVCLQWLKRV QNVSVLSICD FREDWQTWDC PKTQEHIEES RTGWAGGFFV STPVFSYDAI SYYKIFSDKD GYKHIHYIKD TVENAIQITS GKWEAINIFR VTQDSLFFSS NEFEEYPGRR NIYRISIGSY PPSKKCVTCH LRKERCQYYT ASFSYAKYY ALVCYGPPIP ISTLHDGRD QEIKILEENK ELENALKNIQ LPKEEIKKLE VDEITLWYKM ILPPQFDRSK KYPLLIQVYG GPCSQSVRSV FAVNWISYLA SKEGMVIALV DGRGTAFQGD KLLYAVYRKL GVEVEDQIT AVRKFIEMGF IDEKRIAIWG WSYGGYVSSL ALASGTGLFK CGIAPVPSS WEYYASVYTE RFMGLPTKDD NLEHYKNSTV MARAEYFRNV DYLLIHGTAD DNVHFQNSAQ IAKALVNAQV DFQAMWYSDQ NHGLSGLSTN HLYTHMTHFL KQCFSLSD<HH HHHH>

General References

Scanlan, M.J. et al. (1994) Proc. Natl. Acad. Sci. USA 91:5657-5661.
Mori, Y. et al., 2004, Oncology. 67 (5-6):411-419.
O'Brien, P. et al., 2008, Biochim Biophys Acta. 1784 (9):1130-1145.

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain