# NKMAXBio We support you, we believe in your research

# Recombinant human DR3/TNFRSF25 protein

Catalog Number: ATGP4017

## **PRODUCT INFORMATION**

## **Expression system**

Baculovirus

#### **Domain**

25-199aa

#### UniProt No.

093038

#### **NCBI Accession No.**

NP 003781.1

#### **Alternative Names**

Tumor necrosis factor receptor superfamily member 25, Apo-3, Apoptosis-inducing receptor AIR, Apoptosis-mediating receptor DR3, Apoptosis-mediating receptor TRAMP, Death receptor 3, Lymphocyte-associated receptor of death, LARD, Protein WSL, Protein WSL-1, APO3, DDR3, DR3, TNFRSF12, WSL, WSL1, TR3

## **PRODUCT SPECIFICATION**

## **Molecular Weight**

46.1kDa (417aa)

#### Concentration

0.25mg/ml (determined by absorbance at 280nm)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

#### **Purity**

> 90% by SDS-PAGE

#### **Endotoxin level**

< 1 EU per 1ug of protein (determined by LAL method)

## **Biological Activity**

Measured by its binding ability in a functional ELISA with Human TL1A/TNFSF15 (CAT# ATGP1049). The ED50 range  $\leq$  5 ug/ml.

#### Tag

hlgG-His-Tag

#### **Application**

SDS-PAGE, Bioactivity

## **Storage Condition**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.



## Recombinant human DR3/TNFRSF25 protein

Catalog Number: ATGP4017

## **BACKGROUND**

#### **Description**

TNFRSF25, also known as Tumor necrosis factor receptor superfamily member 25, is a member of the TNF receptor superfamily that binds to the TNF-like protein TL1A. TNFRSF25 signals are required to exert T helper cell 2 effector function in Th2-polarized CD4 cells and co-stimulate interleukin-13 production by glycosphingolipidactivated NKT cells. Its axis is a novel immune pathway that participates in the pathogenesis of a variety of autoimmune rheumatic diseases. These molecules may be promising therapeutic targets for inflammatory arthritis. Recombinant human TNFRSF25, fused to hlgG-His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

## **Amino acid Sequence**


<ADP>QGGTRSP RCDCAGDFHK KIGLFCCRGC PAGHYLKAPC TEPCGNSTCL VCPQDTFLAW ENHHNSECAR CQACDEQASQ VALENCSAVA DTRCGCKPGW FVECQVSQCV SSSPFYCQPC LDCGALHRHT RLLCSRRDTD CGTCLPGFYE HGDGCVSCPT STLGSCPERC AAVCGWRQ<LE PKSCDKTHTC PPCPAPELLG GPSVFLFPPK PKDTLMISRT PEVTCVVVDV SHEDPEVKFN WYVDGVEVHN AKTKPREEQY NSTYRVVSVL TVLHQDWLNG KEYKCKVSNK ALPAPIEKTI SKAKGQPREP QVYTLPPSRD ELTKNQVSLT CLVKGFYPSD IAVEWESNGQ PENNYKTTPP VLDSDGSFFL YSKLTVDKSR WQQGNVFSCS VMHEALHNHY TQKSLSLSPG KHHHHHH>

## **General References**

Slebioda TJ., et al. (2011) Eur J Immunol. 41:2606-2611. Fang L., et al. (2008) J Exp Medl. 205:1037-1048.

#### **DATA**

#### SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain

## **Biological Activity**

# **Recombinant human DR3/TNFRSF25 protein**

Catalog Number: ATGP4017

3.0 2.5 2.0 1.5

10-1

Human TNFRSF25 (ug/ml)

Human TL1A/TNFSF15 (CAT# ATGP1049) is coated at 10 ug/ml (100 ul/well) can bind Human DR3/TNFRSF25. The ED50 range  $\leq$  5 ug/ml.