

Biotinylated Recombinant Human Serpin B3/SCCA1 (C-His)

Catalog No: BP090

Description	Biotinylated Recombinant Human Serpin B3 is produced by <i>E.coli</i> . The target gene encoding M1-P390 is expressed with a 6His tag at the C terminus.
Expression System	<i>E. coli</i>
Alternative name	Serpin B3; Protein T4-A; Squamous cell carcinoma antigen 1; SCCA-1; serine (or cysteine) proteinase inhibitor, clade B (ovalbumin), member 3; serpin peptidase inhibitor, clade B (ovalbumin), member 3; Squamous cell carcinoma antigen 1; T4-A; SCCA1
Accession No.	P29508
Predicted Molecular Weight	48.1kDa
Apparent Molecular Weight	SCCA-1 protein appeared at 47kDa in a reducing SDS-PAGE gel
Quality Control	Purity: greater than 95% as determined by reducing SDS-PAGE. Endotoxin: less than 0.1 ng/μg (1 EU/μg) as determined by TAL test.
Formulation	20mM Na ₂ HPO ₄ , 20mM NaH ₂ PO ₄ , pH7.4
Reconstitution	It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Storage	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. Always centrifuge tubes before opening. Do not mix by vortex or pipetting.
Background	Serpin B3, also known as squamous cell carcinoma antigen-1 (SCCA-1), is a member of the serpin superfamily of serine protease inhibitors. Serpins are known to inhibit serine proteases and some can inhibit caspases and papain-like cysteine proteases. Serpin B3 is also a member of the subgroup ovalbumin-related serpins which takes part in the regulation of apoptosis, inflammation, angiogenesis and embryogenesis. Serpin B3 is also a tumor marker for squamous cell carcinomas in diverse areas including the cervix, head, neck and lung. Different from typical serpins, serpin B3 inhibits papain-like lysosomal cysteine proteases Cathepsin K, L, and S, while lacking inhibitory activity against serine proteases

SDS-PAGE

