

Recombinant Human FGF Basic

Catalog No: C046

Description	Recombinant Human Fibroblast growth factor 2/Fibroblast Growth Factor Basic is produced by our E.coli expression system and the target gene encoding Gly132-Ser288 is expressed.
Source	E. coli
Alternative name	Fibroblast Growth Factor 2; FGF-2; Basic Fibroblast Growth Factor; bFGF; Heparin-Binding Growth Factor 2; HBGF-2; FGF2; FGFB
Accession No.	P09038
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.0.
Quality Control	Purity: Greater than 95% as determined by reducing SDS-PAGE. Endotoxin: Less than 0.1 ng/µg (1 IEU/µg).
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Storage	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Background	FGF-basic is a member of the Fibroblast Growth Factors (FGFs) family. The family constitutes a large family of proteins involved in many aspects of development including cell proliferation, growth, and differentiation. They act on several cell types to regulate diverse physiologic functions including angiogenesis, cell growth, pattern formation, embryonic development, metabolic regulation, cell migration, neurotrophic effects, and tissue repair. FGF-basic is a non-glycosylated heparin binding growth factor that is expressed in the brain, pituitary, kidney, retina, bone, testis, adrenal gland liver, monocytes, epithelial cells and endothelial cells. FGF- basic signals through FGFR 1b, 1c, 2c, 3c and 4.

SDS-PAGE

