

Recombinant Human PMP2

Catalog No: C238

Description Recombinant Human Myelin P2 protein is produced by our E.coli expression system and the target

gene encoding Met1-Val132 is expressed with a 6His tag at the N-terminus.

Source E.coli

Alternative name Myelin P2 Protein; Peripheral Myelin Protein 2; PMP2

Accession No. P02689

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Quality Control Purity: Greater than 95% as determined by reducing SDS-PAGE.

Endotoxin: Less than 0.1 ng/μg (1 IEU/μg) as determined by LAL test.

Shipping The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

Storage Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Amino Acid Sequence MGSSHHHHHHSSGLVPRGSHMSNKFLGTWKLVSSENFDDYMKALGVGLATRKLGNLAKPTVIISKKG

DIITIRTESTFKNTEISF

KLGQEFEETTADNRKTKSIVTLQRGSLNQVQRWDGKETTIKRKLVNGKMVAECKMKGVVCTRIYEKV

Reconstitution

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze---thaw cycles.

Background

Myelin P2 Protein (PMP2) is a cytoplasmic protein which belongs to the Fatty-acid binding protein (FABP) family of calycin superfamily. PMP2 is a small, basic, and cytoplasmic lipid binding protein of peripheral myelin. PMP2 is found in peripheral nerve myelin and spinal cord myelin, the oligodendrocytes and Schwann cells, respectively. PMP2 may be involved in lipid transport protein in Schwann cells. It may decrease the inhibitory effect of T suppressors in the culture of immune lymph node cells.

SDS-Page



