

Recombinant Human CDKN1B

Catalog No: C288

Description Recombinant Human Interferon Lambda-1 is produced by our Mammalian expression system and

the target gene encoding Gly20-Thr200 is expressed with a 6His at the C-terminus.

Expression System E.coli

Cyclin-Dependent Kinase Inhibitor 1B; Cyclin-Dependent Kinase Inhibitor p27; p27Kip1; CDKN1B;

Alternative name KIP1

Accession No. P46527

Quality Control Purity: greater than 95% as determined by reducing SDS-PAGE.

Endotoxin: less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Reconstitution It is not recommended to reconstitute to a concentration less than 100µg/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Shipping The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

Storage Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples

are stable at < -20°C for 3 months.

Always centrifuge tubes before opening. Do not mix by vortex or pipetting.

Background Cyclin-Dependent Kinase Inhibitor 1B (CDKN1B) is a Kinesin-related motor protein necessary for

mitotic spindle assembly and chromosome segregation. CDKN1B is expressed in all tissues with highest levels observed in skeletal muscle. CDKN1B is a potent inhibitor of Cyclin E-and Cyclin A-CDK2 complexes. CDKN1B forms a complex with Cyclin Type D-CDK4 complexes and is involved in the assembly, stability, and modulation of CCND1-CDK4

complex activation. In addition, CDKN1B acts as an inhibitor or an activator of Cyclin Type D- CDK4

complexes depending on its phosphorylation state and stoichometry.

SDS-PAGE



