


**Recombinant Human CDH6 (C-6His)**

Catalog No: C434

<b>Description</b>	Recombinant Human Cadherin-6 is produced by our Mammalian expression system and the target gene encoding Thr22-Ala615 is expressed with a 6His tag at the C-terminus.			
<b>Source</b>	Human Cells			
<b>Alternative name</b>	Cadherin-6; Kidney Cadherin; K-Cadherin; CDH6			
<b>Accession No.</b>	P55285			
<b>Predicted Molecular Weight</b>	67.1kDa			
<b>AP Molecular Weight</b>	109kDa, reducing conditions.			
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.			
<b>Reconstitution</b>	<p>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.</p> <p>It is not recommended to reconstitute to a concentration less than 100µg/ml.</p> <p>Dissolve the lyophilized protein in distilled water.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>			
<b>Quality Control</b>	<p>Purity: Greater than 95% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.</p>			
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.			
<b>Storage</b>	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.			
<b>Background</b>	Cadherin-6 (CDH6) is a type-II classic cadherin cell-cell adhesion molecules, which are expressed in graded or areal patterns, as well as layer-specific patterns, in the cortical plate. Human Cadherin-6 is synthesized as a 790 aa type I transmembrane glycoprotein that contains a 18 aa signal peptide, a 35 aa propeptide, a 562 aa extracellular region, a 21 aa transmembrane segment, and a 154 aa cytoplasmic domain. There are five cadherin domains of approximately 110 aa each in the extracellular region. Cadherin-6 has high expression in kidney, brain, and cerebellum, and may contribute to the formation of the segmental structure of the early brain, as well as the development of renal proximal tubules. Weak expression is also detected in lung, pancreas, gastric mucosa and cytotrophoblasts. As a classic cadherin, Cadherin-6 will form homodimers and promote intercellular adhesion with itself and, possibly, Cadherin-9 and -14.			
<b>SDS-Page</b>	<table border="0"> <tr> <td style="text-align: center;">kDa</td> <td style="text-align: center;">MK</td> <td style="text-align: center;">R</td> </tr> </table>  <p>MK: Marker</p> <p>R: Sample under reducing conditions</p>	kDa	MK	R
kDa	MK	R		