

## Recombinant Human DLK-1 (C-6His)

Catalog No: C463

Description Recombinant Human Preadipocyte Factor-1/Protein Delta Homolog 1/Fetal Antigen is produced by our

Mammalian expression system and the target gene encoding Ala24-Pro297(Ser260Asn) is expressed

with a 6His tag at the C-terminus.

Source Human Cells

Alternative name Protein Delta Homolog 1; DLK-1; pG2; DLK1; DLK

Accession No. AAH13197.1

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

**Reconstitution** Always centrifuge tubes before opening. Do not mix by vortex or pipetting.

It is not recommended to reconstitute to a concentration less than 100µg/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Quality Control Purity: Greater than 95% as determined by reducing SDS-PAGE.

Endotoxin: Less than 0.1 ng/μg (1 IEU/μg) as determined by LAL test.

**Shipping** The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

Storage Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Background Protein Delta Homolog 1 (DLK-1) is a transmembrane protein which contains a signal peptide, an

extracellular domain with six tandem epidermai growth factor (EGF)-like domains, a single pass transmembrane domain, and a short cytoplasmic tail. It is found within the stromal cells in close contact to the vascular structure of placental villi, yolk sac, fetal liver, adrenal cortex and pancreas and in the beta cells of the islets of Langerhans in the adult pancreas. In addition, it is detected in some forms of neuroendocrine lung tumor tissue. DLK-1 may have a improtant role in neuroendocrine differentiation.

## SDS-Page



