

## Recombinant Human TrkB (C-6His)

Catalog No: C507

|                                   |   |
|-----------------------------------|---|
| <b>Description</b>                | Recombinant Human Neurotrophic Tyrosine Kinase Receptor Type 2 is produced by our Mammalian expression system and the target gene encoding Cys32-His430 is expressed with a 6His tag at the C- terminus.  |
| <b>Expression System</b>          | Human cells   |
| <b>Alternative name</b>           | BDNF/NT-3 Growth Factors Receptor; GP145-TrkB; Trk-B; Neurotrophic Tyrosine Kinase Receptor Type 2; TrkB Tyrosine Kinase; Tropomyosin-Related Kinase B; NTRK2; TRKB   |
| <b>Accession No.</b>              | Q16220  |
| <b>Predicted Molecular Weight</b> | 45.25kDa  |
| <b>Apparent Molecular Weight</b>  | 60-105kDa, reducing conditions.   |
| <b>Quality Control</b>            | Purity: greater than 95% as determined by reducing SDS-PAGE.<br>Endotoxin: less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.   |
| <b>Formulation</b>                | Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2..  |
| <b>Reconstitution</b>             | It is not recommended to reconstitute to a concentration less than 100μg/ml.<br>Dissolve the lyophilized protein in distilled water.<br>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.   |
| <b>Shipping</b>                   | The product is shipped at ambient temperature.<br>Upon receipt, store it immediately at the temperature listed below.   |
| <b>Storage</b>                    | Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.<br>Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.<br>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.  |
| <b>Background</b>                 | The TRK Family of Tyrosine Kinase Receptor consists of 3 members: TrkA, TrkB and TrkC. The three TRK family proteins have different ligand specificities. They connect to different neurotrophins, including NGF, BDNF, NT-3/NT-4/5. TRKA binds NGF, TRKB binds BDNF and NT-3, TRKC binds NT-4/5. At the protein sequence level, human and rat TRKB have greater than 90% sequence identity and the proteins exhibit cross-species activity. TRKB is primarily expressed in the nervous system and it also expression in a wide variety of tissues with low levels. |

### SDS-PAGE

