

## Recombinant Human DPEP1

Catalog No: C819

---

<b>Description</b>	Recombinant Human Dipeptidase 1 is produced by our Mammalian expression system and the target gene encoding Asp17-Ser385 is expressed with a 6His tag at the C-terminus.
<b>Source</b>	Human Cells
<b>Alternative Name</b>	Dipeptidase 1; Dehydropeptidase-I; Microsomal Dipeptidase; Renal Dipeptidase; hRDP; DPEP1; MDP; RDP
<b>Accession No.</b>	P1644
<b>Predicted Molecular Weight</b>	42.1kDa
<b>Apparent Molecular Weight</b>	41kDa, reducing conditions
<b>Formulation</b>	Supplied as a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH7.4.
<b>Reconstitution</b>	It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
<b>Quality Control</b>	Purity: Greater than 95% as determined by reducing SDS-PAGE. Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Storage</b>	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Background</b>	Dipeptidase 1 (DPEP1) is a kidney membrane enzyme that belongs to the peptidase M19 family. DPEP1 is a homodimer and is inhibited by L-penicillamine. DPEP1 hydrolyzes a variety of dipeptides and is implicated in renal metabolism of glutathione and its conjugates. DPEP1 is responsible for hydrolysis of the beta-lactam ring of antibiotics, such as penem and carbapenem. DPEP1 may play an important role in the regulation of leukotriene activity. DPEP1 expression in cancer is significantly higher than that in normal tissue. However, DPEP1 expression decreased with pathological differentiation, lymph-node and distant metastasis.

### SDS-PAGE

