

Recombinant Human SEMA4D (C-6His)

Catalog No: CC80

Description	Recombinant Human Semaphorin 4D is produced by our Mammalian expression system and the target gene encoding Met22-Arg734 is expressed with a 6His tag at the C-terminus.
Source	Human Cells
Alternative name	Semaphorin-4D; A8;BB18; GR3; CD100.
Accession No.	Q92854
Predicted Molecular Weight	80.2kDa
AP Molecular Weight	1122kDa, reducing conditions.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Reconstitution	<p>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.</p> <p>It is not recommended to reconstitute to a concentration less than 100µg/ml.</p> <p>Dissolve the lyophilized protein in distilled water.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>
Quality Control	<p>Purity: Greater than 90% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.</p>
Shipping	<p>The product is shipped at ambient temperature.</p> <p>Upon receipt, store it immediately at the temperature listed below.</p>
Storage	<p>Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.</p> <p>Reconstituted protein solution can be stored at 4-7°C for 2-7 days.</p> <p>Aliquots of reconstituted samples are stable at < -20°C for 3 months.</p>
Background	Semaphorin-4D is also known as A8,BB18, GR3, CD100. Semaphorin-4D belongs to the semaphorin family containing 1 Ig-like C2-type domain, 1 PSI domain and 1 Sema domain. It is the cell surface receptor for PLXN1B and PLXNB2 that plays an important role in cell-cell signaling. It promotes the migration of cerebellar granule cells and of endothelial cells, regulates dendrite and axon branching and morphogenesis. Semaphorin-4D Plays a role in the immune system; Promotes signaling via SRC and PTK2B/PYK2, which then mediates activation of phosphatidylinositol 3-kinase and of the AKT1 signaling cascade.

SDS-Page

