

Recombinant Mouse CXCL2

Catalog No: CF10

Description	Recombinant Mouse C-X-C motif chemokine 2 is produced by our E. coli expression system and the target gene encoding Ala28-Asn100 is expressed.
Source	E. coli
Alternative name	MIP-2; chemokine ligand 2; C-X-C motif chemokine 2; GRO beta; GRO2; GROB; Gro-beta; Growth-regulated protein beta; Macrophage Inflammatory Protein-2-alpha; melanoma growth stimulatory activity beta; cxcl2; MGSA-b; MGSA-beta; MIP2A; MIP2-alpha; SCYB2.
Accession No.	P10889
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Tris, 150mM NaCl, pH8.0.
Reconstitution	<p>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.</p> <p>It is not recommended to reconstitute to a concentration less than 100µg/ml.</p> <p>Dissolve the lyophilized protein in distilled water.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>
Quality Control	<p>Purity: Greater than 95% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.</p>
Shipping	<p>The product is shipped at ambient temperature.</p> <p>Upon receipt, store it immediately at the temperature listed below.</p>
Storage	<p>Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.</p> <p>Reconstituted protein solution can be stored at 4-7°C for 2-7 days.</p> <p>Aliquots of reconstituted samples are stable at < -20°C for 3 months.</p>
Amino Acid Sequence	AVVASELRCQCLKTLPRVDFKNIQSLSVTPPGPHCAQTEVIATLKGGQKVCCLDPEAPLVQKIIQKILNKG KAN
Background	C-X-C motif chemokine 2 (CXCL2, MIP-2) belongs to the intercrine alpha (chemokine Cx) family. It was originally identified as a heparin-binding protein secreted from a murine macrophage cell line in response to endotoxin stimulation. The expression of mouse MIP-2 is stimulated by endotoxin. The mouse MIP-2 shares approximately 63% aa sequence identity with murine KC, another mouse alpha chemokine, which is induced by PDGF. It has been suggested that mouse KC and MIP-2 are the homologs of the human GROs and rat CINC3s. Chemotactic for human polymorphonuclear leukocytes but does not induce chemokinesis or an oxidative burst. The expression of MIP-2 was found to be associated with neutrophil influx in pulmonary inflammation and glomerulonephritis, suggesting that MIP-2 may contribute to the pathogenesis of inflammatory diseases.

SDS-Page

