

Recombinant Mouse PD-L1

Catalog No: CJ88

Description	Recombinant Mouse Programmed Cell Death 1 Ligand 1 is produced by our Mammalian expression system and the target gene encoding Phe19-Thr238 is expressed with a 6His tag at the C-terminus.
Source	Human Cells
Alternative name	Programmed cell death 1 ligand 1Cd274; programmed cell death 1 ligand 1;PD-L1;PDCD1 ligand 1;programmed death ligand 1;B7 homolog 1;B7-H1;CD274
Accession No.	Q9EP73
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Reconstitution	<p>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.</p> <p>It is not recommended to reconstitute to a concentration less than 100µg/ml.</p> <p>Dissolve the lyophilized protein in distilled water.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>
Quality Control	<p>Purity: Greater than 95% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.</p>
Shipping	<p>The product is shipped at ambient temperature.</p> <p>Upon receipt, store it immediately at the temperature listed below.</p>
Storage	<p>Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.</p> <p>Reconstituted protein solution can be stored at 4-7°C for 2-7 days.</p> <p>Aliquots of reconstituted samples are stable at < -20°C for 3 months.</p>
Amino Acid Sequence	<p>FTITAPKDLVVEYGSNVTMECRFPVERELDLLALVYWEKEDEQVIQFVAGEEDLKPQHSNFRGRASL PKDQLLKGNALQITDVKLQDAGVYCCISYGGADYKRITLKVNPYRKINQRISVDPATSEHELICQAE YPEAEVIWTNSDHQPVSGKRSVTTSTRTEGMLLNVTSSLRVNATANDVFYCTFWRSQPGQNHTAEIIPE LPATHPPQNRTVDHHHHHH</p>
Background	<p>Mouse Programmed cell death 1 ligand 1(Cd274 , PD-L1), is a member of the growing B7 family of immune proteins.It involved in the costimulatory signal essential for T-cell proliferation and IFNG production in a PDCD1-independent manner. Interaction with PDCD1 inhibits T-cell proliferation by blocking cell cycle progression and cytokine production.B7-H1 has been identified as one of two ligands for programmed death1 (PD1), a member of the CD28 family of immunoreceptors. B7-H1 is constitutively expressed in several organs such as heart, skeletal muscle B7-H1 expression is upregulated in a small fraction of activated T and B cells and a much larger fraction of activated monocytes. The costimulatory function of B7-H1 is critical for enhancing maturation and differentiation of T-cells in lymphoid organs.B7-H1 expression is also induced in dendritic cells and keratinocytes after IFN gamma stimulation. Interaction of B7-H1 with PD1 results in inhibition of TCR- mediated proliferation and cytokine production. The B7-H1:PD1 pathway is involved in the negative regulation of some immune responses and may play an important role in the regulation of peripheral tolerance.</p>

SDS-Page

