

## Recombinant Human PD-L1 Catalog No: CM33

Description Recombinant Human Programmed Cell Death 1 Ligand 1 is produced by our Mammalian expression

system and the target gene encoding Phe19-Thr239 is expressed with a Flag tag at the C-terminus.

Source Human cells

Alternative name Programmed Cell Death 1 Ligand 1; PD-L1; PDCD1 Ligand 1; Programmed Death Ligand 1; B7

Homolog 1; B7-H1; CD274; B7H1; PDCD1L1; PDCD1LG1; PDL1

Accession No. Q9NZQ7

**Quality Control** 

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB 150mM Nacl pH 7.4.

Purity: Greater than 95% as determined by reducing SDS-PAGE.

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg).

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting.

It is not recommended to reconstitute to a concentration less than 100μg/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Shipping** The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

Storage Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Background CD274, also known as B7-H1 or programmed death ligand 1 (PD-L1), is a 40 kD type I

transmembrane protein and a member of the B7 family within the immunoglobulin receptor superfamily. Programmed death-1 ligand- 1 (PD-L1, CD274, B7-H1) has been identified as the ligand for the immunoinhibitory receptor programmed death-1(PD1/PDCD1) and has been demonstrated to play a role in the regulation of immune responses and peripheral tolerance. By binding to PD1 on activated T-cells and B-cells, PD-L1 may inhibit ongoing T-cell responses by inducing apoptosis and arresting cell-cycle progression. Accordingly, it leads to growth of immunogenic tumor growth by increasing apoptosis of antigen specific T cells and may contribute to immune evasion by cancers. PD-L1 thus is regarded as promising therapeutic target for human autoimmune disease and

malignant cancers.

## **SDS-PAGE**



