

## Recombinant Human MPF

Catalog No: CM51

<b>Description</b>	Recombinant Human Mesothelin is produced by our Mammalian expression system and the target gene encoding Leu37-Arg286 is expressed with a Fc tag at the C-terminus.
<b>Source</b>	Human Cells
<b>Alternative name</b>	Megakaryocyte potentiating factor; mesothelin; Pre-pro-megakaryocyte-potentiating factor; soluble MPF mesothelin related protein;CAK1; MPF; MSLN; SMR; CAK1; CAK1 antigen
<b>Accession No.</b>	Q13421
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.
<b>Quality Control</b>	Purity: Greater than 95% as determined by reducing SDS-PAGE. Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Storage</b>	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Amino Acid Sequence</b>	LAGETGQEAAPLDGVLANPPNISSLSPRQLLGFPCEVSGLSTERVRELAVALAQKNVKLSTEQLRCLA HRLSEPPEDLDALPLDLLFLNPDAFSGPQACTRFFSRITKANVDLLPRGAPERQRLPAALACWGVRG SLLSEADVRLGGLACDLPGRFVAESAELLRLVSCPGPLDQDQQAARAALQGGGPPYGGPSTWS VSTMDALRGLLPVLGQPIIRSIPQGIVAARQRSSRDPSWRQPERVDDIEGRMDPEKSCDKTHTCPPC PAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYN STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALH NHYTQKSLSLSPGK
<b>Background</b>	Mesothelin is a cell surface glycoprotein whose expression is limited to mesothelial cells of the serosa (pleura, pericardium, and peritoneum) and epithelial cells of the trachea, tonsils, fallopian tube, and kidneys. Mesothelin plays an important role in cell survival, proliferation, migration, invasion, tumor progression, and resistance to chemotherapy. The overexpression of mesothelin can activate NF-κB and signal transducer and activator of transcription 3 (Stat3), inhibit apoptotic signaling and TNF-α-induced apoptosis, and accelerate the G1–S transition. Mesothelin is also found overexpressed in various cancers, including malignant mesothelioma, pancreatic or ovarian carcinoma, sarcomas and in some gastrointestinal or pulmonary carcinomas. As a result of its limited expression in normal tissues, mesothelin has been reported as an ideal tumor-associated marker for the development of targeted therapy.

### SDS-Page

