

Recombinant Mouse SIRPA (C-Fc)

Catalog No: CP68

Description	Recombinant Mouse Signal-Regulatory Protein Alpha 1 is produced by our Mammalian expression system and the target gene encoding Lys32-Asn373 is expressed with a Fc tag at the C-terminus.
Expression System	Human cells
Alternative name	Tyrosine-Protein Phosphatase Non-Receptor Type Substrate 1; CD172 Antigen-Like Family Member A; Signal-Regulatory Protein Alpha-1; CD172a; SIRPA; MFR; SHPS1; SIRP
Accession No.	Q6P6I8
Predicted Molecular Weight	64.8kDa
Apparent Molecular Weight	115.125kDa, reducing conditions.
Quality Control	Purity: greater than 95% as determined by reducing SDS-PAGE. Endotoxin: less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Storage	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. Always centrifuge tubes before opening. Do not mix by vortex or pipetting.
Background	Mouse Signal Regulatory Protein α (SIRPα) is a type I transmembrane glycoprotein. It contains two Ig-like C1-type domains and one Ig-like V-type domain. Mouse SIRP alpha ECD shares 61%, 75%, 62%, 61%, and 59% aa sequence identity with human, rat, equine, bovine, and porcine SIRP alpha, respectively. SIRPα can express in various tissues, mainly on brain and myeloid cells, including macrophages, neutrophils, dendritic and Langerhans cells. It also can detect in neurons, smooth muscle and endothelial cells. SIRPA is an immunoglobulin-like cell surface receptor for CD47. SIRP α acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. SIRP α shows adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. SIRP α engagement generally produces a negative regulatory signal; it may mediate negative regulation of phagocytosis, mast cell activation and dendritic cell activation.

SDS-PAGE

