

Recombinant Mouse CD244 (C-6His)

Catalog No: CS28

Description	Recombinant Mouse Natural killer cell receptor 2B4 is produced by our Mammalian expression system and the target gene encoding Gln20-Asn221 is expressed with a 6His tag at the C-terminus.
Source	Human Cells
Alternative name	Natural killer cell receptor 2B4; NK cell type I receptor protein 2B4; NKR2B4; SLAM family member 4; SLAMF4; Signaling lymphocytic activation molecule 4; CD244
Accession No.	Q07763
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Quality Control	Purity: Greater than 95% as determined by reducing SDS-PAGE. Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Storage	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Background	Natural killer cell receptor 2B4 (2B4/CD244) is a 66 kDa type I transmembrane glycoprotein in the SLAM subgroup of the CD2 protein family. SLAM family proteins have an extracellular domain (ECD) with two or four Ig-like domains and at least two cytoplasmic immunoreceptor tyrosine-based switch motifs (ITSMs). 2B4 interacts with CD48, while other SLAM family proteins interact in a homophilic manner. The mouse 2B4 cDNA encodes a 397 amino acid (aa) precursor that includes a 19 aa signal sequence, a 207 aa ECD with one Ig-like V-type and one C2-type Ig-like domain, a 21 aa transmembrane segment, and a 150 aa cytoplasmic domain with four ITSMs. Within the ECD, mouse 2B4 shares 46% and 68% aa sequence identity with human and rat 2B4, respectively. 2B4/CD48 signaling cooperates with other receptor systems to either promote or inhibit NK and CD8+ T cell activation. The inhibitory activities are distinct from those of MHC I restricted inhibitory NK cell receptors. Ligation of 2B4 with antibodies or CD48 constructs can directly trigger inhibitory signaling or disrupt an inhibitory interaction, leading to cellular activation. 2B4 can also induce signaling through CD48.

