

Recombinant Human CD24

Catalog No: CS99

Description	Recombinant Human Signal Transducer CD24 is produced by our Mammalian expression system and the target gene encoding Ser27-Gly59 is expressed with a Fc tag at the C-terminus.
Source	Baculovirus
Alternative name	Signal transducer CD24; Small cell lung carcinoma cluster 4 antigen; CD24; CD24A
Accession No.	P25063
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.
Quality Control	Purity: Greater than 95% as determined by reducing SDS-PAGE. Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Storage	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Amino Acid Sequence	SETTTGTSSNSSQSTSNTGLAPNPTNATTKAAGIEGRMDPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGSFFLYSKLTVDKSRWQQGNVFSCSVM HEALHNHYTQKSLSLSPGK
Background	Signal Transducer CD24 is a heavily and variably glycosylated GPI-linked sialoprotein. Human CD24 is expressed on B lineage cells and granulocytes, on epithelial, neuronal, and muscle cells, and on a range of tumor cells. CD24 expression is regulated during lineage development and with the activation of various cell types. Antibody crosslinking of CD24 enhances the induction of apoptosis in B and T lymphocytes which contributes to negative selection and the induction of immune tolerance. CD24 on antigen presenting cells cooperates with B7 molecules in the costimulation of T cells. CD24 associates in cis with Siglec10 and with the danger- associated molecules HMGB1, HSP70, or HSP90 which are released from necrotic or damaged cells. Formation of these ternary complexes fills a protective role: the resulting Siglec10 signaling inhibits inflammatory responses that are otherwise induced by extracellular DAMPs.

SDS-Page

