

Human GM-CSF / CSF2 HEK293 Cell Lysate (WB positive control)

Catalog Number: 10015-HNAHL



Sino Biological Inc.
Biological Solution Specialist

GM-CSF Transfected / Overexpression Cell Lysate Product Information

Expressed Host:	HEK293 Cells
Products Description:	Human Cell lysate that Human GM-CSF / CSF2 transfected / overexpressed for Western blot (WB) positive control. The whole cell lysate is provided in 1X Sample Buffer (1X modified RIPA buffer+1X SDS loading buffer).
Sequence information:	A DNA sequence encoding human GMCSF (NP_000749.2) (Met1-Glu144) was expressed.
Predicted N Terminal:	Ala 18
Molecule Mass:	The recombinant human GMCSF consists of 127 amino acids and predicts a molecular mass of 14.5 kDa. As a result of glycosylation, it migrates as an approximately 17-25 kDa band in SDS-PAGE under reducing conditions.
Species:	Human

GM-CSF Transfected / Overexpression Cell Lysate Usage Guide

Preparation Method:	Cell lysate was prepared by homogenization in ice-cold modified RIPA Lysis Buffer with cocktail of protease inhibitors (Sigma). Cell debris was removed by centrifugation. Protein concentration was determined by Bradford assay (Bio-Rad protein assay, Microplate Standard assay). The cell lysate was boiled for 5 min in 1 x SDS loading buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis Buffer:	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF.
Quality Control Testing:	12.5% SDS-PAGE Stained with Coomassie Blue after protein purification.
Stability:	Samples are stable for up to twelve months from date of receipt.
Recommend Usage:	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boil for 2-5 min. 3. Store the lyophilized cell lysate at 4°C. After re-dissolution, recommend to aliquot it into smaller quantities and store at -80°C.
Storage Buffer:	1 X Sample Buffer (1 X modified RIPA buffer+1 X SDS loading buffer).
Storage Instruction:	Store at 4°C. After re-dissolution, aliquot and store at -80°C.
Application notes:	Western blot (WB): Use at an assay dependent dilution. Other Applications: Not tested. Optimal dilutions/concentrations should be determined by the end user.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 ● Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 ● Tel:+86-400-890-9989 ● <http://www.sinobiological.com>