

Human Glypican 3 / GPC3 / OCI-5 Protein (His Tag)

Catalog Number: 10088-H08H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

DGSX; Glypican 3; GTR2-2; MXR7; OCI-5; SDYS; SGB; SGBS; SGBS1

Protein Construction:

A DNA sequence encoding the human GPC3 (P51654-1) (Met 1-His 559) was expressed, with a C-terminal polyhistidine tag.

Source: Human

Expression Host: HEK293 Cells

QC Testing

Purity: > 87 % as determined by SDS-PAGE

Bio Activity:

Measured by its binding ability in a functional ELISA
• Immobilized human GPC3 at 5 µg/ml (100 µl/well) can bind human bFGF / FGF2 with a linear range of 0.156-2.5 ng/ml.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Gln 25 & Ser 359

Molecular Mass:

The secreted recombinant human GPC3 comprises 546 amino acids and has a predicted molecular mass of 62 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Glypican-3, also known as Intestinal protein OCI-5, GPC3, and OCI5, is a member of the glypican family. It belongs to the glypican family and is highly expressed in lung, liver, and kidney. It is a heparan sulfate proteoglycan, which is overexpressed in various neoplasms such as hepatocellular carcinoma, malignant melanoma, and testicular yolk sac tumor, and plays an important role in cell growth and differentiation. GPC3 function is tissue dependent. In some tissues, GPC3 acts as a tumor suppressor gene, whereas in others, it acts as an oncofetal protein. Studies have shown that GPC3 is a reliable marker for hepatocellular carcinoma. The sensitivity and specificity exceeds both alpha-fetoprotein and hepatocyte-paraffin1. GPC3 immunohistochemistry can aid in the differentiation of testicular germ cell tumors, being expressed in all yolk sac tumors but not in seminomas. GPC3 expression has also been identified in some squamous cell carcinomas of the lung and clear cell carcinomas of the ovary. The role of GPC3 in melanomas is still controversial. Thus, Glypican-3 is currently regarded as a tumor marker and potential target for immunotherapy.

References

Kandil DH, et al. (2009) Glypican-3: a novel diagnostic marker for hepatocellular carcinoma and more. Adv Anat Pathol. 16(2): 125-9.
Maeda D, et al. (2009) Glypican-3 expression in clear cell adenocarcinoma of the ovary. Mod Pathol. 22(6): 824-32.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>