

PIK3CG/PI3K Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 100994-T40

GENERAL INFORMATION

Immunogen:	A synthetic peptide corresponding to the C-terminus of the Human PIK3CG/PI3K
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the C-terminus of the Human PIK3CG/PI3K, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human, Mouse
Formulation:	0.2 µm filtered solution in PBS
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

APPLICATIONS

Applications:	WB, IHC-P
---------------	-----------

RECOMMENDED CONCENTRATION

IHC-P	IHC-P: 1:1000-1:5000
Western Blot	WB: 1:500-1:1000


Please Note: Optimal concentrations/dilutions should be determined by the end user.

PIK3CG/PI3K Antibody, Rabbit PAb, Antigen Affinity Purified




Sino Biological
Biological Solution Specialist


Catalog Number: 100994-T40




Immunohistochemical staining PIK3CG in human liver with rabbit polyclonal antibody (1:2500, formalin-fixed paraffin embedded sections).




Immunohistochemical staining PIK3CG in human kidney with rabbit polyclonal antibody (1:2500, formalin-fixed paraffin embedded sections).



Immunohistochemical staining PIK3CG in mouse liver with rabbit polyclonal antibody (1:2500, formalin-fixed paraffin embedded sections).



Immunohistochemical staining CK14 in rat skin with rabbit polyclonal antibody (1:2500, formalin-fixed paraffin embedded sections).



Anti-PIK3CG rabbit polyclonal antibody at 1:500 dilution

Lane A: Jurkat Whole Cell Lysate

Lysates/proteins at 30 µg per lane.

Secondary

Goat Anti- Rabbit IgG H&L (Dylight 800) at 1/10000 dilution.

Developed using the Odyssey technique.

Performed under reducing conditions.

Predicted band size: 126 kDa

Observed band size: 126 kDa