
VEGFR1 / FLT-1 Antibody, Mouse MAb






| GENERAL INFORMATION | |
|---------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Immunogen: | Recombinant Human VEGFR1 / FLT-1 protein (Catalog#10136-H08H) |
| Preparation | This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human VEGFR1 / FLT-1; Catalog#10136-H08H; P17948-1; Met 1-Asn 756). The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography. |
| lg Type: | Mouse IgG2a |
| Clone ID: | 5A4D2 |
| Specificity: | Human VEGFR1 / FLT-1 |
| | No cross-reactivity in ELISA with Human VEGFR2 Human VEGFR3 293 cell lysate |
| Formulation: | 0.2 µm filtered solution in PBS |
| Storage: | This antibody can be stored at $2^{\circ}C-8^{\circ}C$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at $-20^{\circ}C$ to $-80^{\circ}C$. Preservative-Free. Avoid repeated freeze-thaw cycles. |
| Alternative Names: | FLT,FLT-1,VEGFR-1,VEGFR1 |
| APPLICATIONS | |
| Applications: | ELISA,ICC/IF |
| RECOMMENDED CONCENTRATION | |
| ICC/IF | ICC/IF: 1:20-1:100 |
| ELISA | ELISA: 1:1000-1:2000 This antibody can be used at 1:1000-1:2000 with the appropriate secondary reagents to detect Human VEGFR1. |

Please Note: Optimal concentrations/dilutions should be determined by the end user.

VEGFR1 / FLT-1 Antibody, Mouse MAb

Catalog Number: 10136-MM03





Immunofluorescence staining of Human VEGFR1 in Hela cells. Cells were fixed with 4% PFA, blocked with 10% serum, and incubated with Mouse anti-Human VEGFR1 monoclonal antibody (dilution ratio 1:60) at 37°C 1 hour. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-mouse IgG secondary antibody. Positive staining was localized to plasma membrane.