
PAX2 Antibody, Rabbit PAb, Antigen Affinity Purified






GENERAL INFORMATION	
Immunogen:	A synthetic peptide corresponding to the C-terminus of the Human PAX2
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the C-terminus of the Human PAX2, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human, Mouse
	Reacts with Mouse
Formulation:	0.2 µm filtered solution in PBS
Storage:	This antibody can be stored at $2^{\circ}\text{C}-8^{\circ}\text{C}$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C . Preservative-Free. Avoid repeated freeze-thaw cycles.
Alternative Names:	RP11-179B2.1,PAPRS
APPLICATIONS	
Applications:	WB,IHC-P,ICC/IF
RECOMMENDED CONCENTRATION	
IHC-P	IHC-P: 1:2500-1:10000
ICC/IF	ICC/IF: 1:300-1:10000
Western Blot	WB: 1:500-1:2000

Please Note: Optimal concentrations/dilutions should be determined by the end user.


PAX2 Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 101696-T42







Immunochemical staining PAX2 in human kidney with rabbit polyclonal antibody (1:5000, formalin-fixed paraffin embedded sections).




Immunochemical staining PAX2 in human renal carcinoma with rabbit polyclonal antibody (1:5000, formalin-fixed paraffin embedded sections).



Immunochemical staining PAX2 in mouse kidney with rabbit polyclonal antibody (1:5000, formalin-fixed paraffin embedded sections).



Immunofluorescence staining of PAX2 in A549 cells. Cells were fixed with 4% PFA, permeabilzed with 0.1% Triton X-100 in PBS,blocked with 10% serum, and incubated with rabbit anti-Human PAX2 polyclonal antibody (dilution ratio 1:1000) at 4°C overnight. Then cells were stained with the Alexa Fluor\$488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to nucleus.



Anti-PAX2 rabbit polyclonal antibody at 1:500 dilution

Lane A: K562 Whole Cell Lysate Lane B: 293T Whole Cell Lysate Lane C: MCF7 Whole Cell Lysate

Lysates/proteins at 30 µg per lane.

Secondary

Goat Anti-Rabbit InG (H+I

Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique.Performed under reducing conditions.

Predicted band size:44 kDa Observed band size:44 kDa