

MAF Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 102568-T36

GENERAL INFORMATION

Immunogen:	A synthetic peptide corresponding to the N-terminus of the Human MAF
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the N-terminus of the Human MAF, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human Mouse, Rat (Species predicted to react based on 100% sequence homology)
Formulation:	0.2 µm filtered solution in PBS
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

APPLICATIONS

Applications:	WB,IP
----------------------	-------

RECOMMENDED CONCENTRATION

Western Blot	WB: 1:500-1:1000
Immunoprecipitation	IP: 0.2-1 µL/mg of lysate


Please Note: Optimal concentrations/dilutions should be determined by the end user.

MAF Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 102568-T36



Anti-MAF rabbit polyclonal antibody at 1:500 dilution
Lane A: PC12 Whole Cell Lysate


Lysates/proteins at 30 µg per lane.

Secondary

Goat Anti- Rabbit IgG H&L (Dylight 800) at 1/10000 dilution.

Developed using the Odyssey technique.
Performed under reducing conditions.

Predicted band size: 42 kDa
Observed band size: 47 kDa



MAF was immunoprecipitated using:
Lane A: 0.5 mg 293T Whole Cell Lysate

1 µL anti-MAF rabbit polyclonal antibody and
15 µl of 50 % Protein G agarose.

Primary antibody:

Anti-MAF rabbit polyclonal antibody, at 1:500 dilution

Secondary antibody:

Dylight 800-labeled antibody to rabbit IgG (H+L), at 1:5000 dilution

Developed using the odssey technique.
Performed under reducing conditions.

Predicted band size: 41 kDa
Observed band size: 41 kDa