

FAM98A Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 103080-T36

GENERAL INFORMATION

Immunogen:	A synthetic peptide corresponding to the C-terminus of the Human FAM98A
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the C-terminus of the Human FAM98A , and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human Mouse, Rat (Species predicted to react based on 100% sequence homology)
Formulation:	0.2 µm filtered solution in PBS
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

APPLICATIONS

Applications:	WB,IP
----------------------	-------

RECOMMENDED CONCENTRATION

Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP: 4-8 µL/mg of lysate


Please Note: Optimal concentrations/dilutions should be determined by the end user.

FAM98A Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 103080-T36




Anti-FAM98A rabbit polyclonal antibody at
1:500 dilution
Lane A: MCF7 Whole Cell Lysate

Lysates/proteins at 30 µg per lane.
Secondary
Goat Anti- Rabbit IgG H&L (Dylight 800) at
1/10000 dilution.

Developed using the Odyssey technique.
Performed under reducing conditions.

Predicted band size: 55 kDa
Observed band size: 55 kDa



FAM98A was immunoprecipitated using:
Lane A: 0.5 mg 293T Whole Cell Lysate
Lane B: 0.5 mg HepG2 Whole Cell Lysate

4 µL anti-FAM98A rabbit polyclonal antibody
and 15 µL of 50 % Protein G agarose.

Primary antibody:
Anti-FAM98A rabbit polyclonal antibody, at
1:100 dilution

Secondary antibody:
Clean-Blot® IP Detection Reagent (HRP) at
1:500 dilution

Developed using the DAB staining technique.
Performed under reducing conditions.

Predicted band size: 57 kDa
Observed band size: 57 kDa