

# Human Angiopoietin-2 / ANG2 Protein (His Tag), Biotinylated

Catalog Number: 10691-H08S-B



Sino Biological  
Biological Solution Specialist

## General Information

### Gene Name Synonym:

AGPT2; ANG2; Angiopoietin-2

### Protein Construction:

A DNA sequence encoding the human ANGPT2 (NP\_001138.1) (Met1-Phe496) was expressed with a polyhistidine tag at the C-terminus. The purified protein was biotinylated in vitro.


**Source:** Human

**Expression Host:** CHO Stable Cells

## QC Testing

**Purity:** > 95 % as determined by SDS-PAGE.

### SDS-PAGE:



## Protein Description

Angiopoietin-2 (ANG 2, or ANGPT2), is a member of the ANG family, which plays an important role in angiogenesis during the development and growth of human cancers. Both ANGPT-1 and ANGPT-2 appear to bind to the tyrosine kinase receptor, Tie-2, found primarily on the luminal surface of endothelial cells. ANG-2's role in angiogenesis generally is considered as an antagonist for ANG1, inhibiting ANG1-promoted Tie2 signaling, which is critical for blood vessel maturation and stabilization. ANG-2 modulates angiogenesis in a cooperative manner with another important angiogenic factor, vascular endothelial growth factor A. Genetic studies have revealed that ANG-2 also is critical in lymphangiogenesis during development. ANG-2 has multiple physiologic effects that regulate vascular tone, hormone secretion, tissue growth and neural activity. Several reports indicate that ANG-2 can induce neovascularization in experimental systems due to the expression of different growth factors such as angiopoietin 2, vascular endothelial factor, and its receptor, fibroblast growth factor, platelet derived growth factor, transforming growth factor beta and epidermal growth factor. In addition, ANG-2 is strongly expressed in the vasculature of many tumors and it has been suggested that ANG-2 may act synergistically with other cytokines such as vascular endothelial growth factor to promote tumor-associated angiogenesis and tumor progression.

## Molecular Mass:

The recombinant human ANGPT2 consists of 489 amino acids and predicts a molecular mass of 56.4 kDa.

### Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

## Usage Guide

### Storage:

Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

**Avoid repeated freeze-thaw cycles.**

## References

- 1.Thomas M, et al. (2009) The role of the Angiopoietins in vascular morphogenesis. *Angiogenesis.* 12(2): 125-37.
- 2.Hu B, et al. (2009) Angiopoietin-2: development of inhibitors for cancer therapy. *Curr Oncol Rep.* 11(2): 111-6.
- 3.Fiedler U, et al. (2006) Angiopoietins: a link between angiogenesis and inflammation. *Trends Immunol.* 27: 552-8.

### Reconstitution:

Detailed reconstitution instructions are sent along with the products.

**Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.**

**For US Customer: Fax: 267-657-0217**

● **Tel: 215-583-7898**

**Global Customer: Fax :+86-10-5862-8288**

● **Tel:+86-400-890-9989** ● <http://www.sinobiological.com>