
SNAP-25 Antibody, Rabbit PAb, Antigen Affinity Purified






GENERAL INFORMATION	
Immunogen:	A synthetic peptide corresponding to the N-terminus of the Human SNAP-25
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the N-terminus of the Human SNAP-25, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human, Mouse, Rat Bovine, Chick, Pongo abelii (Species predicted to react based on 100% sequence homology)
Formulation:	0.2 µm filtered solution in PBS
Storage:	This antibody can be stored at $2^{\circ}\text{C-8}^{\circ}\text{C}$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C . Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.
APPLICATIONS	
Applications:	WB, IHC-P
RECOMMENDED CONCENTRATION	
IHC-P	IHC-P: 1:500-1:2000
Western Blot	WB: 1:500-1:2000

Please Note: Optimal concentrations/dilutions should be determined by the end user.

SNAP-25 Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 107191-T40






Immunochemical staining of human SNAP25 in human brain with rabbit polyclonal antibody at 1:1000 dilution, formalin-fixed paraffin embedded sections.



Immunochemical staining of human SNAP25 in human liver with rabbit polyclonal antibody at 1:1000 dilution, formalin-fixed paraffin embedded sections.




Immunochemical staining of human SNAP25 in mouse brain with rabbit polyclonal antibody at 1:1000 dilution, formalin-fixed paraffin embedded sections.



Immunochemical staining of human SNAP25 in mouse liver with rabbit polyclonal antibody at 1:1000 dilution, formalin-fixed paraffin embedded sections.



Immunochemical staining of human SNAP25 in rat brain with rabbit polyclonal antibody at 1:1000 dilution, formalin-fixed paraffin embedded sections.




Immunochemical staining of human SNAP25 in rat liver with rabbit polyclonal antibody at 1:1000 dilution, formalin-fixed paraffin embedded sections.

SNAP-25 Antibody, Rabbit PAb, Antigen Affinity Purified







Anti-SNAP25 rabbit polyclonal antibody at 1:500 dilution Lane A: SH-SY5Y Whole Cell Lysate

Lysates/proteins at 30 µg per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique. Performed under reducing conditions.

Predicted band size:23 kDa Observed band size:26 kDa