

Human DDR1 Kinase / MCK10 / CD167 Protein (aa 444-913, His & GST Tag)

Catalog Number: 10730-H20B1



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

CAK; CD167; DDR; EDDR1; HGK2; MCK10; NEP; NTRK4; PTK3; PTK3A; RTK6; TRKE

Protein Construction:

A DNA sequence encoding the human DDR1 (Q08345-1) (Arg444-Val913) was fused with the N-terminal polyhistidine-tagged GST tag at the N-terminus.

Source: Human

Expression Host: Baculovirus-Insect Cells

QC Testing

Purity: > 80 % as determined by SDS-PAGE

Bio Activity:

The specific activity was determined to be 2.75 nmol/min/mg using synthetic AXLtide peptide(CKKSRGDYMTMQIG) as substrate.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Met

Molecular Mass:

The recombinant human DDR1 /GST chimera consists of 707 amino acids and has a calculated molecular mass of 80 kDa. The recombinant protein migrates as an approximately 80 kDa band in SDS-PAGE under reducing conditions.

Formulation:

Supplied as sterile 20mM Tris, 500mM NaCl, pH 7.4, 10% glycerol, 3mM DTT

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Discoidin domain receptor family, member 1 (DDR1), also known as or CD167a (cluster of differentiation 167a), and Mammary carcinoma kinase 10 (MCK10), belongs to a subfamily of tyrosine kinase receptors with an extracellular domain homologous to Dictyostellium discoideum protein discoidin 1. Receptor tyrosine kinases play a key role in the communication of cells with their microenvironment. These kinases are involved in the regulation of cell growth, differentiation and metabolism. Expression of DDR1/MCK10/CD167 is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, it has been shown to be significantly overexpressed in several human tumors. DDR1/MCK10/CD167 plays an important role in regulating attachment to collagen, chemotaxis, proliferation, and MMP production in smooth muscle cells. DDR1 functions in a feedforward loop to increase p53 levels and at least some of its effectors. Inhibition of DDR1 function resulted in strikingly increased apoptosis of wild-type p53-containing cells in response to genotoxic stress through a caspase-dependent pathway.

References

1. Hou G, *et al.* (2001) The discoidin domain receptor tyrosine kinase DDR1 in arterial wound repair. *J Clin Invest.* 107(6): 727-35.
2. Ongusaha PP, *et al.* (2003) p53 induction and activation of DDR1 kinase counteract p53-mediated apoptosis and influence p53 regulation through a positive feedback loop. *EMBO J.* 22(6): 1289-301.
3. Jönsson M, *et al.* (2001) Repression of Wnt-5a impairs DDR1 phosphorylation and modifies adhesion and migration of mammary cells. *J Cell Sci.* 114(11): 2043-53.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>