

Human CD113 / Nectin-3 / PVRL3 Protein (His Tag)

Catalog Number: 10852-H08H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

CD113; CDw113; DKFZp566B0846; FLJ90624; nectin-3; PPR3; PRR3; PVRL3; PVRR3

Protein Construction:

A DNA sequence encoding the extracellular domain of human PVRL3 (NP_056295.1) (Met 1-Asp 400) was expressed, with a polyhistidine tag at the C-terminus.

Source: Human

Expression Host: HEK293 Cells

QC Testing

Purity: > 98 % as determined by SDS-PAGE

Bio Activity:

Measured by its binding ability in a functional ELISA . 1. Immobilized recombinant human PVRL3 at 1 µg/ml (100 µl/well) can bind biotinylated Nectin-1 with a linear range of 6.4-800 ng/ml . 2. Immobilized recombinant human PVRL3 at 1 µg/ml (100 µl/well) can bind Nectin-1/ Fc chimera with a linear range of 0.156-5 ng/ml.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Gly 58

Molecular Mass:

The secreted recombinant human PVRL3 comprises 354 amino acids and has a predicted molecular mass of 39.3 kDa. As a result of glycosylation, rhPVRL3 migrates as an approximately 65-75 kDa band in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Poliovirus receptor-related 3 (PVRL3), also known as Nectin-3 and CD113, is a member of the nectin family. PVRL3/Nectin-3 is an 83 kDa, type I transmembrane glycoprotein. Its precursor is 549 amino acids (aa) in length and contains an extended signal sequence of 57 aa, an extracellular domain (ECD) of 347 aa, a transmembrane segment of 21 aa, and a cytoplasmic region of 124 aa. Nectin-3 has three splicing variants, nectin-3alpha (biggest), -3beta (middle), and -3gamma (smallest). It is predominantly expressed in testis and placenta as well as in various cell lines, including epithelial cell lines. PVRL3/Nectin-3 plays a role in cell-cell adhesion through heterophilic trans-interactions with nectin-like proteins or nectins, such as trans-interaction with PVRL2/Nectin-2 at Sertoli-spermatid junctions. PVRL3/Nectin-3 is thus involved in the formation of cell-cell junctions, including adherens junctions and synapses. It has been shown to induce endocytosis-mediated down-regulation of PVR from the cell surface, resulting in reduction of cell movement and proliferation.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>