Human ENTPD3 / NTPDase3 / CD39L3 Protein (His Tag)

Catalog Number: 10909-H08B



General Information

Gene Name Synonym:

CD39L3; HB6; NTPDase-3

Protein Construction:

A DNA sequence encoding the human ENTPD3 (O75355) extracellular domain (Gln 44-Pro 485) was fused with a polyhistidine tag at the C-terminus and a signal peptide at the N-terminus.

Source: Human

Expression Host: Baculovirus-Insect Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

Measured by its ability to hydrolyze the 5'-phosphate group from the substrate adenosine-5'-triphosphate (ATP). The orthophosphate product is measured by a Malachite Green Phosphate Detection Kit (Catalog # DY996). The specific activity is >70,000 pmol/min/µg.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt $% 10^{\circ}$ at -70 $^{\circ}\mathrm{C}$

Predicted N terminal: Gln 44

Molecular Mass:

The recombinant human ENTPD3 (aa44-485) consists of 452 amino acids and predicts a molecular mass of 51 kDa. It migrates as an approximately 50 KDa band in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 7.4, 10% gly

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide


Storage:

Store it under sterile conditions at -20 $^\circ\!\!\!C$ to -80 $^\circ\!\!\!C$ upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.



References

1.Fausther M, *et al.* (2010) Cystic fibrosis remodels the regulation of purinergic signaling by NTPDase1 (CD39) and NTPDase3. Am J Physiol Lung Cell Mol Physiol. 298(6): 804-18. 2.Vlajkovic SM, *et al.* (2006) Noise-induced up-regulation of NTPDase3 expression in the rat cochlea: Implications for auditory transmission and cochlear protection. Brain Res. 1104(1): 55-63. 3.Ivanenkov w, *et al.* (2005) Characterization of disulfide bonds in human nucleoside triphosphate diphosphohydrolase 3 (NTPDase3): implications for NTPDase structural modeling. Biochemistry. 44 (25): 8998-9012.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 2

Global Customer: Fax :+86-10-5862-8288

Tel: 215-583-7898

Tel:+86-400-890-9989 •

http://www.sinobiological.com

SDS-PAGE: