

Rhesus PCSK9 / NARC1 Protein (His Tag)

Catalog Number: 11054-K08H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

PCSK9

Protein Construction:

A DNA sequence encoding the Rhesus PCSK9 (NP_001106130.1) (Met 1-Gln 692) precursor was expressed, fused with a polyhistidine tag at the C-terminus.

Source: Rhesus

Expression Host: HEK293 Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

1. Measured by its ability to bind biotinylated human LDLR in a functional ELISA . 2. Measured by its ability to bind biotinylated mouse LDLR in a functional ELISA.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Gln 31

Molecular Mass:

The recombinant Rhesus PCSK9 consists of 673 amino acids and predicts a molecular mass of 72.7 kDa. As a result of glycosylation and proteolytic digestion, it migrates as doublet with apparent molecular mass of 20 kDa and 62 kDa corresponding to the pro domain and mature form respectively in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Proprotein convertase subtilisin/kexin type 9 (PCSK9), also known as NARC1 (neural apoptosis regulated convertase), which is a newly identified human secretory subtilase belonging to the proteinase K subfamily of the secretory subtilase family. PCSK9 protein is an enzyme which in humans is encoded by the PCSK9 gene with orthologs found across many species. It is expressed in neuroepithelioma, colon carcinoma, hepatic and pancreatic cell lines, and in Schwann cells. PCSK9 protein is highly expressed in the liver and regulates low density lipoprotein receptor (LDLR) protein levels. Inhibition of PCSK9 protein function is currently being explored as a means of lowering cholesterol levels. Thereby, PCSK9 protein is regarded as a new strategy to treat hypercholesterolemia. PCSK9 protein contributes to cholesterol homeostasis and may have a role in the differentiation of cortical neurons. References

References

- 1.Sseidah, N.G. et al., 2003, Proc. Natl. Acad. Sci. USA. 100: 928-933.
- 2.Beyer, T.P. et al., 2007, J. Lipid. Res. 48: 1488-1498
- 3.Shan, L. et al., 2008, Biochem. Biophys. Res. Commun. 375: 69-73.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>