Influenza A H5N1 (A/bar-headed goose/Qinghai/14/2008) Hemagglutinin / HA HEK293 Cell Lysate (WB positive control)

Catalog Number: 11059-V08H2L



Hemagglutinin / HA Transfected / Overexpression Cell Lysate Product Information

Expressed Host: HEK293 Cells

Products Description: Human Cell lysate that Influenza A H5N1 (A/bar-headed goose/Qinghai/14/2008) Hemagglutinin / HA

transfected / overexpressed for Western blot (WB) positive control. The whole cell lysate is provided in

1X Sample Buffer (1X modified RIPA buffer+1X SDS loading buffer).

Sequence information: A DNA sequence encoding the extracellular domain of Influenza A virus (A/bar-headed

goose/Qinghai/14/2008 (H5N1)) (ACL28277.1) HA (Met 1-Gln 531) (Native, HA1+HA2, cleaved) was

expressed with a C-terminal polyhistidine tag.

Predicted N Terminal: Asp 17

Molecule Mass: The secreted recombinant hemagglutinin of Influenza A virus (A/bar-headed goose/Qinghai/14/2008

(H5N1) comprises 526 amino acids with the predicted molecular mass of 60 kDa. As a result of glycosylation, it migrates as an approximately 70 kDa band in Non-reduced SDS-PAGE, and two bands

(28, 50 kDa) in reduced SDS-PAGE.

Species: H5N1

Hemagglutinin / HA Transfected / Overexpression Cell Lysate Usage Guide

Preparation Method: Cell lysate was prepared by homogenization in ice-cold modified RIPA Lysis Buffer with cocktail of

protease inhibitors (Sigma). Cell debris was removed by centrifugation. Protein concentration was determined by Bradford assay (Bio-Rad protein assay, Microplate Standard assay). The cell lysate was boiled for 5 min in 1 x SDS loading buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium

dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.

Lysis Buffer: Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1%

SDS, 1% Sodium deoxycholate, 1mM PMSF.

Quality Control Testing: 12.5% SDS-PAGE Stained with Coomassie Blue after protein purification.

Stability: Samples are stable for up to twelve months from date of receipt.

Recommend Usage:

1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.

2. Re-dissolve the pellet using 200µL pure water and boil for 2-5 min.

3. Store the lyophilized cell lysate at 4°C. After re-dissolution, recommend to aliquot it into smaller

quantities and store at -80°C.

Storage Buffer: 1 X Sample Buffer (1 X modified RIPA buffer+1 X SDS loading buffer).

Storage Instruction: Store at 4°C. After re-dissolution, aliquot and store at -80°C.

Application notes: Western blot (WB): Use at an assay dependent dilution.

Other Applications: Not tested.

Optimal dilutions/concentrations should be determined by the end user.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • http://www.sinobiological.com