

# Human R-Spondin 1 / RSPO1 Protein (His Tag)

Catalog Number: 11083-H08H



Sino Biological  
Biological Solution Specialist

## General Information

### Gene Name Synonym:

CRISTIN3; R-Spondin 1; RSPO

### Protein Construction:

A DNA sequence encoding the human RSPO1 (NP\_001033722.1) (Met 1-Ala 263) was expressed fused with a polyhistidine tag at the C-terminus.

**Source:** Human

**Expression Host:** HEK293 Cells

## QC Testing

**Purity:** > 95 % as determined by SDS-PAGE

### Bio Activity:

1. Measured by its binding ability in a functional ELISA. Immobilized human RSPO1 at 20 µg/ml (100 µl/well) can bind human LIMP2 with a linear range of 32-800 ng/ml.
2. Measured by its binding ability in a functional ELISA. Immobilized human RSPO1 at 20 µg/ml (100 µl/well) can bind mouse CD36 with a linear range of 6.4-800 ng/ml.
3. Measured by its ability to induce activation of βcatenin response in a Topflash Luciferase assay using HEK293T human embryonic kidney cells. The ED<sub>50</sub> for this effect is typically 0.1-0.9 µg/mL in the presence of 5 ng/mL recombinant mouse Wnt3a.

### Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

**Predicted N terminal:** Ser 21

### Molecular Mass:

The secreted recombinant human RSPO1 comprises 254 amino acids with a predicted molecular mass of 28.2 kDa. As a result of glycosylation, rhRSPO1 migrates as an approximately 42 kDa band in SDS-PAGE under reducing conditions.

### Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

## Usage Guide

### Stability & Storage:

Samples are stable for twelve months from date of receipt at -20°C to -80°C.


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

**Avoid repeated freeze-thaw cycles.**

### Reconstitution:

Detailed reconstitution instructions are sent along with the products.

## SDS-PAGE:



## Protein Description

RSPO1 gene is a member of the R-spondin family. It encodes RSPO1 which is known as a secreted activator protein with two cysteine-rich, furin-like domains and one thrombospondin type 1 domain. In mice, RSPO1 induces the rapid onset of crypt cell proliferation and increases intestinal epithelial healing, providing a protective effect against chemotherapy-induced adverse effects. This protein is an activator of the beta-catenin signaling cascade, leading to TCF-dependent gene activation. RSPO1 acts both in the canonical Wnt/beta-catenin-dependent pathway and in non-canonical Wnt signaling pathway, probably by acting as an inhibitor of ZNRF3, an important regulator of the Wnt signaling pathway. It also acts as a ligand for frizzled FZD8 and LRP6.

## References

1. Kamata T, et al. (2004) R-spondin, a novel gene with thrombospondin type 1 domain, was expressed in the dorsal neural tube and affected in Wnts mutants. *Biochim Biophys Acta*. 1676(1):51-62.
2. Ota T, et al. (2004) Complete sequencing and characterization of 21,243 full-length human cDNAs. *Nat Genet*. 36(1):40-5.
3. Strausberg RL, et al. (2003) Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. *Proc Natl Acad Sci*. 99(26):16899-903.

**For Research Use Only. Not for use in diagnostic or therapeutic procedures.**

**Tel: +86-400-890-9989 (Global), +1-215-583-7898 (USA), +49(0)6196 9678656 (Europe)**

**Website: <http://www.sinobiological.com>**