

Human R-Spondin 1 / RSPO1 Protein

Catalog Number: 11083-HNAS



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

CRISTIN3; R-Spondin 1; RSPO

Protein Construction:

A DNA sequence encoding the human RSPO1 (NP_001033722.1) (Met 1-Ala 263) was expressed and purified.

Source: Human

Expression Host: CHO Stable Cells

QC Testing

Purity: (53.2+43.4) % as determined by SDS-PAGE

Bio Activity:

Measured by its ability to induce activation of β catenin response in a Topflash Luciferase assay using HEK293T human embryonic kidney cells. The ED_{50} for this effect is typically 20-120 ng/mL in the presence of 5 ng/mL recombinant mouse Wnt3a.

Endotoxin:

< 1.0 EU per μ g of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal: Ser 21

Molecular Mass:

The recombinant human RSPO1 consists of 243 amino acids and has a predicted molecular mass of 26.8 kDa. As a result of glycosylation, the apparent molecular mass of it is approximately 40 and 31 kDa in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, PH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

RSPO1 gene is a member of the R-spondin family. It encodes RSPO1 which is known as a secreted activator protein with two cystein-rich, furin-like domains and one thrombospondin type 1 domain. In mice, RSPO1 induces the rapid onset of crypt cell proliferation and increases intestinal epithelial healing, providing a protective effect against chemotherapy-induced adverse effects. This protein is an activator of the beta-catenin signaling cascade, leading to TCF-dependent gene activation. RSPO1 acts both in the canonical Wnt/beta-catenin-dependent pathway and in non-canonical Wnt signaling pathway, probably by acting as an inhibitor of ZNRF3, an important regulator of the Wnt signaling pathway. It also acts as a ligand for frizzled FZD8 and LRP6.

References

1. Kamata T, *et al.* (2004) R-spondin, a novel gene with thrombospondin type 1 domain, was expressed in the dorsal neural tube and affected in Wnts mutants. *Biochim Biophys Acta.* 1676(1):51-62.
2. Ota T, *et al.* (2004) Complete sequencing and characterization of 21,243 full-length human cDNAs. *Nat Genet.* 36(1):40-5.
3. Strausberg RL, *et al.* (2003) Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. *Proc Natl Acad Sci.* 99(26):16899-903.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>