

Human CTLA4 / CD152 Protein (His & Fc Tag)



Sino Biological
Biological Solution Specialist

Catalog Number: 11159-H03H

General Information

Gene Name Synonym:

ALPS5; CD; CD152; CELIAC3; CTLA-4; GRD4; GSE; IDDM12

Protein Construction:

A DNA sequence encoding the human CTLA4 (NP_005205.2) extracellular domain (Met 1-Phe 162) was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.

Source: Human

Expression Host: HEK293 Cells

QC Testing

Purity: > 92 % as determined by SDS-PAGE

Bio-Activity:

Measured by its ability to inhibit IL-2 secretion by stimulated Jurkat human acute T cell leukemia cells. The ED₅₀ for this effect is 0.2-1 µg/mL when stimulated with 1 µg/mL Recombinant Human B7-1/CD80.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Ala 37

Molecular Mass:

The recombinant human CTLA4/Fc is a disulfide-linked homodimer. The reduced monomer consists of 374 amino acids and has a predicted molecular mass of 41.6 kDa. As a result of glycosylation, the apparent molecular mass of rh CTLA4/Fc monomer is approximately 50-55 kDa in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:

Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:


Detailed reconstitution instructions are sent along with the products.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>

SDS-PAGE:



Protein Description

Cytotoxic T-lymphocyte protein 4, also known as CTLA4 and CD152, is a single-pass type I membrane protein and a member of the immunoglobulin superfamily. It is the second member of the CD28 receptor family. The ligands or counterreceptors for these two proteins are the B7 family members, CD80 (B7-1) and CD86 (B7-2). CTLA4 transmits an inhibitory signal to T cells, whereas CD28 transmits a stimulatory signal. Intracellular CTLA4 is also found in regulatory T cells and may play an important role in their functions. CD152 or cytotoxic T lymphocyte antigen-4 (CTLA-4) is an essential receptor involved in the negative regulation of T cell activation. Because of its profound inhibitory role, CD152 has been considered a sound susceptible candidate in autoimmunity and a persuasive target for cancer immunotherapy. In particular, recent evidence suggests that CD152 is also important in the homeostasis and function of a population of suppressive cells, termed regulatory T cells (Treg).

References

- 1.Slavik JM, *et al.* (1999) CD28/CTLA-4 and CD80/CD86 families: signaling and function. *Immunol Res.* 19(1): 1-24.
- 2.Holmberg D, *et al.* (2005) CTLA-4 (CD152) and its involvement in autoimmune disease. *Autoimmunity.* 38(3): 225-33.
- 3.Chin LT, *et al.* (2008) Immune intervention with monoclonal antibodies targeting CD152 (CTLA-4) for autoimmune and malignant diseases. *Chang Gung Med J.* 31(1): 1-15.