

Human B7-H3 Protein (His & AVI Tag), Biotinylated

Catalog Number: 11188-H27H-B



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

4Ig-B7-H3; B7-H3; B7H3; B7RP-2

Protein Construction:

A DNA sequence encoding the human CD276 (Q5ZPR3) (Met1-Thr461) was expressed with a c-terminal polyhistidine tagged AVI tag at the C-terminus. The expressed protein was biotinylated in vivo by the Biotin-Protein ligase (BirA enzyme) which is co-expressed.

Source: Human

Expression Host: Human Cells

QC Testing

Biotin/Protein Ratio:

0.5-1 as determined by the HABA assay.

Purity: > 95 % as determined by SDS-PAGE.

Endotoxin:

< 1.0 EU per µg protein as determined by the LAL method.

Predicted N terminal: Leu 29

Molecular Mass:

The recombinant human CD276 consists of 459 amino acids and predicts a molecular mass of 49.8 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Stability & Storage:

Samples are stable for twelve months from date of receipt at -20°C to -80°C.


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

B7-H3 is a member of the B7 family of immune regulatory ligands that is thought to attenuate peripheral immune responses through co-inhibition. It plays an important role in adaptive immune responses, and was shown to either promote or inhibit T-cell responses in various experimental systems. B7-H3 may play an important role in muscle-immune interactions, providing further evidence of the active role of muscle cells in local immunoregulatory processes. B7-H3 is a novel protein structurally related to the B7 family of ligands by the presence of a single set of immunoglobulin-V-like and immunoglobulin-C-like (VC) domains. Previous studies have correlated its overexpression with poor prognosis and decreased tumor-infiltrating lymphocytes in various carcinomas including uterine endometrioid carcinomas, and mounting evidence supports an immuno-inhibitory role in ovarian cancer prognosis. Recently, B7-H3 expression has been reported in several human cancers indicating an additional function of B7-H3 as a regulator of antitumor immunity.

References

1. Suh WK, *et al.* (2004) The immune regulatory protein B7-H3 promotes osteoblast differentiation and bone mineralization. *Proc Natl Acad Sci U S A.* 101(35): 12969-73.
2. Waschbisch A, *et al.* (2008) Human muscle cells express the costimulatory molecule B7-H3, which modulates muscle-immune interactions. *Arthritis Rheum.* 58(11): 3600-8.
3. Loos M, *et al.* (2010) B7-h3 and its role in antitumor immunity. *Clin Dev Immunol.* 2010: 683875.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

Tel: +86-400-890-9989 (Global), +1-215-583-7898 (USA), +49(0)6196 9678656 (Europe)

Website: <http://www.sinobiological.com>