# Human CADM3 / NECL1 / IGSF4B Protein (His Tag)

Catalog Number: 11214-H08H



## **General Information**

#### Gene Name Synonym:

BlgR; IGSF4B; Necl-1; NECL1; synCAM3; TSLL1

#### **Protein Construction:**

A DNA sequence encoding the human CADM3 isoform 2 (short isoform) (NP\_001120645.1) extracellular domain (Met 1-His 330) was expressed, with a polyhistidine tag at the C-terminus.

Source: Human

Expression Host: HEK293 Cells

**QC** Testing

Purity: > 95 % as determined by SDS-PAGE

## **Bio Activity:**

Measured by the ability of the immobilized protein to support the adhesion of C6 Rat brain glial cells. Human CADM3 immobilized (0.8  $\mu$ g/ml, 100  $\mu$ l/well) will mediate >30% C6 cell adhesion.

#### **Endotoxin:**

< 1.0 EU per µg of the protein as determined by the LAL method

#### Stability:

Samples are stable for up to twelve months from date of receipt at -70  $^{\circ}\mathrm{C}$ 

Predicted N terminal: Asn 25

#### **Molecular Mass:**

The recombinant human CADM3 consists of 317 amino acids and predictes a molecular mass of 35.1 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rh CADM3 is approximately 39-41 kDa due to glycosylation.

## Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

#### **Usage Guide**

### Storage:


Store it under sterile conditions at  $-20\,^\circ\mathbb{C}$  to  $-80\,^\circ\mathbb{C}$  upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

## Avoid repeated freeze-thaw cycles.

#### Reconstitution:

Detailed reconstitution instructions are sent along with the products.

#### SDS-PAGE:



## **Protein Description**

Cell Adhesion Molecules (CAMs) are proteins located on the cell surface involved with the binding with other cells or with the extracellular matrix (ECM) in the process called cell adhesion. These proteins are typically transmembrane receptors and are composed of three domains: an intracellular domain that interacts with the cytoskeleton, a transmembrane domain, and an extracellular domain that interacts either with other CAMs of the same kind (homophilic binding) or with other CAMs or the extracellular matrix (heterophilic binding). Cell adhesion molecule 3, also known as Immunoglobulin superfamily member 4B, CADM3, and NECL1, is a neural tissue-specific immunoglobulin-like cell-cell adhesion molecule which has Ca(2+)-independent homo- or heterophilic cell-cell adhesion activity and plays an important role in the formation of synapses, axon bundles and myelinated axons. Isoform 1 of CADM3 is expressed mainly in adult and fetal brain. Isoform 2 of CADM3 is highly expressed in adult brain and weakly expressed in placenta. In brain, Isoform 2 is highly expressed in cerebellum. CADM3 is involved in the cell-cell adhesion. It has both calcium-independent homophilic cell-cell adhesion activity and calciumindependent heterophilic cell-cell adhesion activity with IGSF4, PVRL1 and PVRL3. The interaction with EPB41L1 may regulate structure or function of cell-cell junctions. CADM3 may act as a tumor suppressor in glioma and loss of it in glioma may be caused by histone deacetylation.

#### References

1.Dong X, et al. (2006) Crystal structure of the V domain of human Nectin-like molecule-1/Syncam3/Tsll1/lgsf4b, a neural tissue-specific immunoglobulin-like cell-cell adhesion molecule. J Biol Chem. 281(15): 10610-7. 2.Gao J, et al. (2008) Nectin-like molecule 1 is a glycoprotein with a single N-glycosylation site at N290KS which influences its adhesion activity. Biochim Biophys Acta. 1778(6): 1429-35. 3.Gao J, et al. (2009) Loss of NECL1, a novel tumor suppressor, can be restored in glioma by HDAC inhibitor-Trichostatin A through Sp1 binding site. Glia. 57(9): 989-99.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 
■ Tel:+86-400-890-9989 
■ http://www.sinobiological.com