
ICOS Ligand / B7-H2 Antibody, Mouse MAb







GENERAL INFORMATION	
Immunogen:	Recombinant Human ICOS Ligand / B7-H2 Protein (Catalog#11559-H08H)
Preparation	This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human ICOS Ligand / B7-H2 (rh ICOS Ligand / B7-H2; Catalog#11559-H08H; NP_056074.1; Met1-Ser258). The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.
lg Type:	Mouse IgG1
Clone ID:	01
Specificity:	Human ICOS Ligand / B7-H2
Formulation:	0.2 µm filtered solution in PBS
Storage:	This antibody can be stored at $2^{\circ}\text{C-8}^{\circ}\text{C}$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C . Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.
APPLICATIONS	
Applications:	FCM
RECOMMENDED CONCENTRATION	
Flow Cytometry	FCM: 1:25-1:100

Please Note: Optimal concentrations/dilutions should be determined by the end user.

ICOS Ligand / B7-H2 Antibody, Mouse MAb







Flow cytometric analysis of Human ICOSLG(CD275) expression on RPMI8226 cells. Cells were stained with purified anti-Human ICOSLG(CD275), then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.

Flow cytometry was performed on a BD FACSCalibur flow cytometry system. Please refer to www.sinobiological.com/Flow-Cytometry-FACS-Protocols-a-750.html for technical protocols.