

Human CD200R1L / CD200R2 / CD200RLa Protein (His Tag)

Catalog Number: 11620-H08H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

CD200R2; CD200RLa

Protein Construction:

A DNA sequence encoding the human CD200R1L (AAT00538.1) extracellular domain (Met1-Leu239) was expressed, fused with a polyhistidine tag at the C-terminus.

Source: Human

Expression Host: HEK293 Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Gly 24

Molecular Mass:

The recombinant human CD200R1L consists of 227 amino acids and predicts a molecular mass of 25.2 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rh CD200R1L is approximately 45-60 kDa due to glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Cell surface glycoprotein CD200 receptor 2, also known as Cell surface glycoprotein CD200 receptor 1-like, Cell surface glycoprotein OX2 receptor 2, CD200 receptor-like 2, CD200R1a, CD200R1L and CD200R2, is a single-pass type I membrane protein which belongs to the CD200R family. CD200R1L / CD200R2. It contains one Ig-like C2-type (immunoglobulin-like) domain and one Ig-like V-type (immunoglobulin-like) domain. CD200 is a transmembrane protein delivering immunoregulatory signals after engagement of CD200R. A family of CD200Rs exist (CD200R1, CD200R2, CD200R3, CD200R4) with different tissue expression and functional activity. In the presence of anti-CD200R2 / CD200R3 monoclonal antibodies (mAbs), bone-marrow cells cultured in the presence of (interleukin [IL]-4+granulocyte-macrophage colony-stimulating factor) differentiate into dendritic cells (DCs), which induce CD4+CD25+ Treg. Interaction between the relatively ubiquitously expressed molecule CD200 and one of its receptors, CD200R1, resulted in direct suppression of alloreactivity, engagement of alternate receptors led instead to altered differentiation of dendritic cells (DCs) from marrow precursors, which could in turn foster development of Foxp3(+) regulatory T cells. Unlike anti-CD200R1, anti-CD200R2 both promotes development of DCs with capacity to induce Treg and directly augments thymocyte production of Treg.

References

1. Wright G.J., *et al.*, (2003), Characterization of the CD200 receptor family in mice and humans and their interactions with CD200. *J. Immunol.* 171:3034-3046.
2. Gorczynski R.M., *et al.*, (2004), Structural and functional heterogeneity in the CD200R family of immunoregulatory molecules and their expression at the feto-maternal interface. *Am. J. Reprod. Immunol.* 52:147-163.
3. Muzny D.M., *et al.*, (2006), The DNA sequence, annotation and analysis of human chromosome 3. *Nature* 440:1194-1198.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>