

Human IL17RC Protein (aa 1-454, His Tag)

Catalog Number: 11747-H08B



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

IL17-RL; IL17RL

Protein Construction:

A DNA sequence encoding the full length of human IL17RC isoform 3 (NP_116121.2) (Met 1-Ala 454) was fused with a polyhistidine tag at the carboxy-terminus.

Source: Human

Expression Host: Baculovirus-Insect Cells

QC Testing

Purity: > 90 % as determined by SDS-PAGE

Bio Activity:

1. Measured by its ability to bind with recombinant human IL17A-His (Cat:12047-H07B) in a functional ELISA. 2. Measured by its ability to bind with recombinant human 17A (Cat:12047-HNAE) in a functional ELISA.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Leu 21

Molecular Mass:

The recombinant human IL17RC (aa 1-454) consists of 444 amino acids and predicts a molecular mass of 49.6 kDa. The apparent molecular mass of the protein is approximately 60 kDa in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 7.4, 10% gly

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



References

1.Allen W, *et al.* (2010) IL-17RC: A partner in IL-17 signaling and beyond. *Semin Immunopathol.* 32(1): 33-42. 2.You ZB, *et al.* (2007) Differential Expression of IL-17RC Isoforms in Androgen-Dependent and Androgen-Independent Prostate Cancers. *Neoplasia.* 9(6): 464-70. 3.Zrioual S, *et al.* (2008) IL-17RA and IL-17RC receptors are essential for IL-17A-induced ELR+ CXC chemokine expression in synoviocytes and are overexpressed in rheumatoid blood. *J Immunol.* 180(1): 655-63.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>