

Human FES Kinase / Feline sarcoma oncogene Protein (His & GST Tag)

Catalog Number: 12214-H20B



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

FPS

Protein Construction:

A DNA sequence encoding the human FES (P07332-1) (Met 1-Arg 822) was fused with the N-terminal polyhistidine-tagged GST tag at the N-terminus.

Source: Human

Expression Host: Baculovirus-Insect Cells

QC Testing

Purity: > 75 % as determined by SDS-PAGE

Bio Activity:

The specific activity was determined to be 200 nmol/min/mg using Poly(Glu:Tyr) 4:1 as substrate.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Met

Molecular Mass:

The recombinant human FES/GST chimera consists of 1060 amino acids and has a calculated molecular mass of 121 kDa. It migrates as an approximately 110 kDa band in SDS-PAGE under reducing conditions.

Formulation:

Supplied as sterile 20mM Tris, 500mM NaCl, pH 7.4, 10% glycerol

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Proto-oncogene tyrosine-protein kinase Fes/Fps, also known as Proto-oncogene c-Fes, Proto-oncogene c-Fps, Feline sarcoma oncogene, FES and FPS, is a protein which contains oneFCH domain, oneprotein kinase domain and oneSH2 domain. FES is a non-receptor protein tyrosine kinase expressed in hematopoietic progenitors and differentiated myeloid cells. FES is observed in the nuclear, granular and plasma membrane fractions of primary human neutrophils and the myeloid leukemia cell line, HL-60. The nuclear localization is confirmed by immunocytochemistry of neutrophils. FES has been implicated in granulocyte-macrophage colony-stimulating factor (GM-CSF), interleukin-3 (IL-3) and erythropoietin signal transduction. FES has tyrosine-specific protein kinase activity and that activity is required for maintenance of cellular transformation. FES is also involved in normal hematopoiesis. Its chromosomal location has linked it to a specific translocation event identified in patients with acute promyelocytic leukemia.

References

- 1.Bowden DW, et al.,1991, Nucleic acids Res 19 (15): 4311.
- 2.Yates,K.E. et al., 1995, Oncogene. 10 (6):1239-42.
- 3.Jücker, M, et al.,1997, J. Biol. Chem. 272 (4): 2104-9.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>