Human Resistin / ADSF / RETN Protein (Fc Tag)

Catalog Number: 13055-H02H



General Information

Gene Name Synonym:

ADSF; FIZZ3; RETN1; RSTN; XCP1

Protein Construction:

A DNA sequence encoding the human RETN (NP_065148.1) (Ser17-Pro108) was expressed with the Fc region of human IgG1 at the C-terminus.

Source: Human

Expression Host: HEK293 Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE.

Endotoxin:

< 1.0 EU per μg protein as determined by the LAL method.

Stability:

Samples are stable for up to twelve months from date of receipt $\,$ at -70 $\,$ $^{\circ}$ C

Predicted N terminal: Glu

Molecular Mass:

The recombinant human RETN consists 352 amino acids and predicts a molecular mass of 38.2 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20 $^\circ\!\mathrm{C}$ to -80 $^\circ\!\mathrm{C}$ upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



References

1.Verma S, et al. (2003) Resistin promotes endothelial cell activation: further evidence of adipokine-endothelial interaction. Circulation. 108(6):736-40. 2.Reilly MP, et al. (2005) Resistin is an inflammatory marker of atherosclerosis in humans. Circulation. 111(7):932-9. 3.Steppan CM, et al. (2002) Resistin and obesity-associated insulin resistance. Trends Endocrinol Metab. 13(1):18-23.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • http://www.sinobiological.com