

Human OX-40L / TNFSF4 / CD252 Protein (Fc Tag)

Catalog Number: 13127-H04H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

CD134L; CD252; GP34; OX-40L; OX4OL; TXGP1

Protein Construction:

A DNA sequence encoding the human TNFSF4 (NP_003317.1) (Gln51-Leu183) was expressed with the Fc region of mouse IgG1 at the N-terminus.

Source: Human

Expression Host: HEK293 Cells

QC Testing

Purity: > 90 % as determined by SDS-PAGE.

Bio Activity:

Measured by its binding ability in a functional ELISA. Immobilized recombinant Human TNFRSF4-His (Cat:10481-H08H) at 10 µg/ml (100 µl/well) can bind human S4-mFc3-TNFSF4 (Cat:13127-H04H) with a linear range of 1.28-20 µg/ml.

Endotoxin:

< 1.0 EU per µg protein as determined by the LAL method.

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Asp

Molecular Mass:

The recombinant human TNFSF4 consists of 369 amino acids and predicts a molecular mass of 42.1 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

OX-4L, also known as TNFSF4 and CD252, is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. OX-4L is an important costimulatory molecule that plays a crucial role in the regulation of T-cell-mediated immunity. The interaction of TNFSF4-TNFSF4 is involved in the pathogenesis of multiple autoimmune and inflammatory diseases such as systemic lupus erythematosus (SLE), carotid artery disease and cancer. OX-4L is a ligand for receptor TNFRSF4/OX4. It is found to play a role in T cell antigen-presenting cell (APC) interactions. In surface Ig- and CD4-stimulated B cells, this cytokine along with CD7 has been shown to provide CD28-independent costimulatory signals to T cells. This protein and its receptor are reported to directly mediate adhesion of activated T cells to vascular endothelial cells.

References

1.Lei W. et al., 2012, Ann Acad Med Singapore. 41 (5): 200-4. 2.Lee YH. et al., 2012, Hum Immunol. 73 (10): 1050-4. 3.Weiguang Y. et al., 2012, PLoS One. 7 (8): e41277.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 ● Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 ● Tel:+86-400-890-9989 ● <http://www.sinobiological.com>