## **ENO2** Antibody, Rabbit PAb, Antigen Affinity Purified






| GENERAL INFORMATION       |   |
|---------------------------|---|
| Immunogen:                | Recombinant Human ENO2 Protein (Catalog#13219-H08E)   |
| Preparation               | Produced in rabbits immunized with purified, recombinant Human ENO2 (rh ENO2; Catalog#13219-H08E; P09104; Met1-Leu434). ENO2 specific IgG was purified by Human ENO2 affinity chromatography.   |
| Ig Type:                  | Rabbit IgG  |
| Specificity:              | Human ENO2  |
| Formulation:              | 0.2 µm filtered solution in PBS   |
| Storage:                  | This antibody can be stored at $2^{\circ}\text{C}-8^{\circ}\text{C}$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at $-20^{\circ}\text{C}$ to $-80^{\circ}\text{C}$ . Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles. |
| APPLICATIONS              |   |
| Applications:             | IHC-P,IP  |
| RECOMMENDED CONCENTRATION |   |
| IHC-P                     | IHC-P: 1:10000-1:20000  |
| Immunoprecipitation       | IP: 1-4 μL/mg of lysate   |

Please Note: Optimal concentrations/dilutions should be determined by the end user.

## **ENO2** Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 13219-T44






Immunochemical staining of human ENO2 in human pancreas with rabbit polyclonal antibody (1:20000, formalin-fixed paraffin embedded sections).



Immunochemical staining of human ENO2 in human brain with rabbit polyclonal antibody (1:20000, formalin-fixed paraffin embedded sections).



ENO2 was immunoprecipitated using: Lane A:0.5 mg Hela Whole Cell Lysate

4  $\mu L$  anti-ENO2 rabbit polyclonal antibody and 15  $\mu l$  of 50 % Protein G agarose.

Primary antibody: Anti-ENO2 rabbit polyclonal antibody,at 1:100 dilution

Secondary antibody: Clean-Blotô IP Detection Reagent (HRP) at 1:500 dilution

Developed using the DAB staining technique. Performed under reducing conditions.

Predicted band size: 47 kDa Observed band size: 47 kDa