## Actopaxin Antibody, Rabbit PAb, Antigen Affinity Purified






GENERAL INFORMATION	
Immunogen:	Recombinant Human Actopaxin protein (Catalog#13919-H09E)
Preparation	Produced in rabbits immunized with purified, recombinant Human Actopaxin (rh Actopaxin; Catalog#13919-H09E; Q9NVD7-1; Met1-Glu372). Actopaxin specific IgG was purified by Human Actopaxin affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human Actopaxin
Formulation:	0.2 µm filtered solution in PBS
Storage:	This antibody can be stored at $2^{\circ}\text{C-8}^{\circ}\text{C}$ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at $-20^{\circ}\text{C}$ to $-80^{\circ}\text{C}$ . Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.
APPLICATIONS	
Applications:	WB,ELISA,IP
RECOMMENDED CONCENTRATION	
Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP: 1-4 µL/mg of lysate
ELISA	ELISA: 1:5000-1:10000 This antibody can be used at 1:5000-1:10000 with the appropriate secondary reagents to detect Human Actopaxin.
Oleman Nata o Ontire all account out in a fall things about the data wait and be the	

Please Note: Optimal concentrations/dilutions should be determined by the end user.

## Actopaxin Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: 13919-T52





Anti-PARVA rabbit polyclonal antibody at 1:500 dilution

Lane A: HepG2 Whole Cell Lysate Lane B: A549 Whole Cell Lysate Lane C: MCF7 Whole Cell Lysate

Lysates/proteins at 30 µg per lane. Secondary Goat Anti-Rabbit IgG H&L (Dylight800) at 1/10000 dilution.

Developed using the Odyssey technique. Performed under reducing conditions.

Predicted band size:47 kDa Observed band size:48 kDa (We are unsure as to the identity of these extra bands.)



PARVA was immunoprecipitated using: Lane A:0.5 mg HepG2 Whole Cell Lysate

 $2~\mu L$  anti-PARVA rabbit polyclonal antibody and 15  $\mu l$  of ~50~%~ Protein G agarose.

Primary antibody: Anti-PARVA rabbit polyclonal antibody,at 1:200 dilution

Secondary antibody: Clean-Blotô IP Detection Reagent (HRP) at 1:1000 dilution

Developed using the DAB staining technique. Performed under reducing conditions.

Predicted band size: 50 kDa Observed band size: 50 kDa