

Human ECD / Ecdysoneless homolog Protein (His & GST Tag)

Catalog Number: 14311-H20B



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

ECD; GCR2; HSGT1

Protein Construction:

A DNA sequence encoding the human ECD (O95905-1) (Met1-Thr621) was expressed with the N-terminal polyhistidine-tagged GST tag at the N-terminus.

Source: Human

Expression Host: Baculovirus-Insect Cells

QC Testing

Purity: > 90 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Met

Molecular Mass:

The recombinant human ECD/GST chimera consists of 881 amino acids and has a calculated molecular mass of 100.6 kDa. The recombinant protein migrates as an approximately 88-108 kDa band in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile 20mM Tris, 500mM NaCl, 10% glycerol, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

ECD, also known as ecdysoneless homolog, belongs to the SGT1 family. It is highly expressed in muscle and heart. ECD is a novel promoter of mammalian cell cycle progression. This function is related to its ability to remove the repressive effects of Rb-family tumor suppressors on E2F transcription factors. It is a novel tumor-promoting factor that is differentially expressed in pancreatic cancer and potentially regulates glucose metabolism within cancer cells. ECD may also be a transcriptional activator required for the expression of glycolytic genes.

References

- 1.Badzek S. et al., 2011, Wien Klin Wochenschr. 123 (23-24): 726-31.
- 2.Zhao X. et al., 2012, Breast Cancer Res Treat. 134 (1): 171-80. 3.Dey P. et al., 2012, Clin Cancer Res. 18 (22): 6188-98.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • **Tel: 215-583-7898**

Global Customer: Fax :+86-10-5862-8288 • **Tel:+86-400-890-9989** • <http://www.sinobiological.com>